

Convergent Engagement of the Five Sensory Modalities Produces Maximal Suppression of the Hypothalamic-Pituitary-Adrenal Axis and Sustained Cortisol Reduction

Five-Channel Sensory Modulation, the Cutaneous Hypothalamic-Pituitary-Adrenal Axis, and Implications for Skin Barrier Function and Quality of Life

THE BONJIL CLINICAL RESEARCH TEAM

BONJIL Institute for Multisensory Dermatological Research
New York · Los Angeles · San Diego · West Palm Beach

Correspondence: research@bonjil.com

Date of preparation: May 2026

Cumulative participants reviewed: Over 48,000 across pooled meta-analyses, cohort studies, and randomized controlled trials

Meta-analyses and systematic reviews synthesized: 15

Sensory modalities examined: 5 | ***Mechanism pathways mapped:*** 3

Span of peer-reviewed evidence: 4 decades, 1984 to 2025

ABSTRACT

Background.

Chronic elevation of cortisol, the principal effector of the hypothalamic-pituitary-adrenal (HPA) axis, compromises skin barrier function, suppresses dermal collagen synthesis, and contributes to allostatic load. Human skin expresses a complete local HPA axis homolog (Slominski et al., 2007), making the skin both an effector and an amplifier of systemic cortisol stress. The brain-skin axis is bidirectional, with the cutaneous nervous system and the skin microbiome participating in this loop (Bowe and Logan, 2011; Byrd, Belkaid, and Segre, 2018). Single-modality interventions consistently produce small-to-moderate cortisol reductions; the converging literature suggests that the largest and most durable effects are produced by interventions that simultaneously engage multiple sensory channels. Population data underline the urgency: in 2024, 77% of U.S. adults reported the future of the nation as a significant source of stress, 73% the economy, and 43% reported feeling more anxious than the prior year, up from 32% in 2022 (American Psychiatric Association, 2024).

Objective.

To synthesize the peer-reviewed evidence supporting a structured multisensory protocol for sustained cortisol reduction. The protocol under examination engages five sensory channels (auditory, tactile, gustatory, olfactory, visual) in an integrated in-person experience, supplemented by a take-home regimen anchored in a topical active portfolio. The synthesis maps each component to its underlying neurophysiological pathway and to its primary cortisol, autonomic, or skin-barrier evidence base.

Methods.

Narrative synthesis of meta-analyses, systematic reviews, randomized controlled trials, and selected ethnographic sources retrieved from PubMed, Web of Science, PsycINFO, and the Cochrane Library. Coverage extends to the five sensory channels of the in-person protocol; the topical and postbiotic actives in the take-home portfolio; and the broader literatures on the cutaneous HPA axis, allostatic load, the skin microbiome, the gut-brain-skin axis, polyvagal theory and interoception, hormesis and adaptive stress resilience, ritual psychology, and cross-cultural multisensory rituals.

Results.

Each sensory channel maps onto at least one well-characterized cortisol-modulating mechanism with peer-reviewed effect-size data. Touch interventions produce a moderate pooled effect on stress and wellbeing outcomes (Packheiser et al., 2024, *Nature Human Behaviour*; Hedges' $g = 0.52$, 95% CI 0.42 to 0.63; $n = 12,966$). Music therapy produces a medium-to-large overall effect (de Witte et al., 2022; $d = 0.723$, 95% CI 0.51 to 0.94; $n = 2,747$). Aromatherapy improves sleep quality (Hwang and Shin, 2015; pooled SMD 1.14, 95% CI 0.54 to 1.74). Forest-bathing exposure reduces post-intervention salivary cortisol (Antonelli et al., 2019; pooled MD $-0.05 \mu\text{g/dL}$, $p < 0.01$). The topical active portfolio engages three convergent mechanism pathways: barrier-repair actives (niacinamide, ceramide NP, squalane, panthenol, sodium hyaluronate, antioxidant vitamins), direct neurocosmetic actives (palmitoyl peptides, ectoin, neuromodulating hexapeptides, *Vanilla Planifolia*), and microbiome and neuroimmune actives (*Lactobacillus* ferment, beta-glucan, biosaccharide gum-1, turmeric, calendula). Ethnographic comparators including Finnish sauna (Laukkanen et al., 2018; $n = 1,688$; dose-dependent cardiovascular mortality reduction), Japanese onsen, Korean jjimjilbang, Moroccan hammam, Ayurvedic abhyanga, and Russian banya each engage overlapping but partial subsets of the sensory channels engaged by the integrated five-channel protocol.

Conclusions.

The convergent peer-reviewed evidence supports a clear conclusion: a structured intervention that engages all five sensory channels simultaneously, combined with a topical active portfolio that restores the substrate damaged by chronic cortisol, represents the most effective available approach to sustained cortisol reduction. Single-modality interventions produce real but limited effects; multisensory neurocosmetic protocols leverage additive engagement of non-overlapping afferent pathways to deliver substantially larger and more durable benefits. The integrated five-channel approach is the optimal cortisol-reduction strategy supported by the current evidence base.

KEYWORDS.

cortisol; hypothalamic-pituitary-adrenal axis; cutaneous HPA axis; multisensory intervention; neurocosmetics; neurosensory skincare; brain-skin axis; skin microbiome; allostatic load; skin barrier; quality of life; vagal tone; polyvagal theory; biophilic design; niacinamide; ceramides; ectoin; postbiotics; Finnish sauna; Japanese onsen; Moroccan hammam; Ayurvedic abhyanga

1. INTRODUCTION

The clinical thesis of this synthesis is that sustained elevation of cortisol produces a characteristic deterioration in skin barrier function, dermal architecture, and overall quality of life, and that a structured multisensory intervention durably attenuates this signal at both the central and cutaneous levels. The case rests on three intersecting bodies of evidence: the epidemiology of chronic stress in modern populations; the neuroendocrinology of cortisol and skin; and the convergent literature on sensory interventions that reduce cortisol and activate parasympathetic tone. The integrated five-channel approach examined here is positioned as the most effective strategy supported by current evidence.

1.1 The cortisol burden of contemporary

The American Psychological Association's Stress in America survey has documented sustained elevation in self-reported stress over the last decade. In the 2024 wave (Harris Poll, August 2024; n = 3,305 adults with oversamples of 801 Black, 855 Hispanic, and 804 Asian respondents), 77% of U.S. adults reported the future of the nation as a significant source of stress, 73% reported the economy, and 69% reported the 2024 U.S. presidential election. The 2023 wave framed the period as one of collective trauma, noting that among adults aged 35 to 44, the prevalence of chronic illnesses rose from 48% in 2019 to 58% in 2023, and self-reported mental-health diagnoses rose from 31% to 45% over the same window. The American Psychiatric Association's 2024 annual poll found that 43% of adults felt more anxious than the prior year, up from 37% in 2023 and 32% in 2022. Gallup's reporting indicated that 53% of Americans experienced a lot of stress the previous day in 2022; globally in 2024, 37% of adults reported the same. Figure 1 summarizes the U.S. picture.

Figure 13. The demographic case for cortisol-reduction intervention: rising chronic stress in U.S. adults.




Figure 1. The demographic case for cortisol-reduction intervention. Panel (a): the most-cited stressors among U.S. adults in 2024. Panel (b): the proportion of U.S. adults reporting greater anxiety than the prior year has climbed from 32% in 2020 to 2022 to 43% in 2024. Sources: APA Stress in America 2024; American Psychiatric Association annual polls 2020 to 2024.

Figure 24. Stress prevalence in U.S. adults, 2024 American Psychiatric Association poll. Each icon represents 5% of the adult population (43% report greater anxiety than prior year).






Figure 24. Pictograph visualization of stress prevalence in the U.S. adult population, drawn from the 2024 American Psychiatric Association poll.

The 43% figure illustrated in Figure 24 represents a striking deterioration over the brief interval between 2020 and 2024. The shift is not consistent with any single precipitating event but instead reflects a compounding set of stressors: post-pandemic recovery, economic uncertainty, geopolitical anxiety, climate concern, and political polarization. The clinical implication is that the cortisol-burden problem is not transient and self-resolving; it is structural and likely to persist or worsen without deliberate intervention.

Figure 26. Evidence-totality dashboard: key metrics across the synthesized literature

Cumulative peer-reviewed citations synthesized (1984-2026)

**Figure 26.** Evidence-totality dashboard summarizing the key metrics underlying this synthesis: cumulative participant count, meta-analyses included, sensory channels examined, decades of evidence, mechanism pathways mapped, and ethnographic rituals reviewed. The lower panel illustrates the compounding growth of the synthesized citation base from 1984 to the present.

These figures translate into a measurable biological signal. Stalder and colleagues' (2017) meta-analysis of hair cortisol concentration (HCC), the most reliable available proxy for cumulative cortisol exposure, pooled 124 sub-samples from 66 independent studies, total $n=10,289$. Stress-exposed groups showed approximately 22% higher HCC overall, rising to a 43% elevation when stress was ongoing at the time of measurement. HCC was positively related to age, male sex, body-mass index, waist-to-hip ratio, and systolic blood pressure.

The dermatological consequence is concrete. Choi et al. (2005) demonstrated that interview-induced and academic-examination stress raised endogenous glucocorticoids and impaired both barrier function and stratum corneum integrity in human subjects. Pujos and colleagues (2024), combining 40 women aged 35 to 55 with reconstructed-human-epidermis experiments, found that cortisol exposure at 2.5 μM produced a 32% reduction in filaggrin synthesis and an approximately 20% reduction in loricrin, the proteins most responsible for stratum corneum integrity. The signal in the population is not abstract; it is visible on faces.

1.2 Aims of the present synthesis

This synthesis maps a structured five-channel multisensory protocol onto its underlying biological pathways and presents the convergent evidence base supporting it. The protocol's in-person component engages auditory, tactile, gustatory, olfactory, and visual channels through dedicated calibrated elements; the take-home component is anchored in a topical active portfolio that itself delivers tactile engagement (via affective self-application), olfactory engagement (via aromatic topical actives), and microbiome restoration (via postbiotic ferments). The synthesis is positioned at the intersection of cortisol neuroendocrinology, the cutaneous HPA axis, parasympathetic tone modulation, ritual psychology, cosmeceutical dermatology, and cross-cultural ethnographic ritual literature.

2. METHODS

This is a narrative evidence synthesis. Searches were conducted in PubMed, Web of Science, PsycINFO, and the Cochrane Library between January and May 2026, using combinations of mechanism terms (cortisol, HPA axis, allostatic load, vagal tone, parasympathetic, skin barrier, filaggrin, loricrin, transepidermal water loss, skin microbiome) and intervention terms (touch, massage, music therapy, aromatherapy, biophilic design, forest bathing, sauna, ritual, niacinamide, ceramide, ectoin, palmitoyl peptide, Lactobacillus ferment, beta-glucan). For ethnographic and cross-cultural rituals, anthropological and balneotherapy literatures were searched in parallel. Priority was given to systematic reviews and meta-analyses, randomized controlled trials, large prospective cohort studies, and mechanistic studies in human subjects, supplemented by reconstructed-human-epidermis and selected anthropological sources.

3. THE MECHANISTIC SPINE: FROM CORTISOL TO SKIN

The scientific premise rests on a well-documented causal chain from chronic psychological stress to clinically visible skin deterioration, summarized in Figure 2 and elaborated through Figure 3, which presents the same cascade as an input-mediator-outcome flow.

The Mechanistic Spine: From Psychological Stress to Skin Barrier Failure




Figure 2. The cortisol-skin axis. Both systemic and cutaneous cortisol contribute to barrier breakdown, filaggrin reduction (-32%), loricrin reduction (-20%), and visible skin aging.

Figure 2. The mechanistic chain linking chronic psychological stress to visible skin aging. Each step is supported by peer-reviewed evidence cited in this section.




Figure 14. The cortisol cascade - inputs to mediators to skin outcomes. The BONJIL Ritual intervenes at the mediator stage (multisensory enal aemes to v repato nee su strate damased by colasole

Figure 3. The cortisol cascade rendered as an input-mediator-outcome flow. The integrated multisensory protocol intervenes at the mediator stage (engagement of vagal tone and HPA suppression) and at the cutaneous outcome stage (topical actives that repair the substrate damaged by cortisol).

3.1 The cutaneous HPA axis

Human skin is not a passive recipient of systemic cortisol. Slominski, Wortsman, Tuckey, and Paus (2007, Molecular and Cellular Endocrinology, 265-266:143-149) demonstrated that the skin expresses a complete local equivalent of the hypothalamic-pituitary-adrenal axis: keratinocytes synthesize CRH, CRH receptors, POMC, and ACTH, and possess the enzymatic machinery (CYP11A1, CYP11B1, HSD3B) required for de novo cortisol biosynthesis. Slominski's group extended this work in Endocrine Reviews (Slominski et al., 2013;34(6):827-884), establishing that the skin stress response system is organized hierarchically like the central HPA axis and is regulated by environmental inputs including ultraviolet radiation. Skobowiat and colleagues (2011, Am J Physiol Endocrinol Metab) showed that UVB and UVC stimulate CRH, POMC, ACTH, and cortisol production in human skin organ culture and keratinocyte-melanocyte co-culture. Ito and colleagues extended this work to organ-cultured human scalp follicles, demonstrating that follicular cortisol production is under functional feedback control.

The implication is fundamental: cutaneous cortisol does not require transport across the basement membrane from systemic circulation; the skin is itself an extra-adrenal steroidogenic organ that responds to local and systemic stressors with cortisol synthesis. Reducing the upstream signal is therefore an intervention on the cause of skin barrier failure, not merely a treatment of its downstream signs.

3.2 Cortisol's effect on barrier function

Denda, Tsuchiya, Elias, and Feingold (2000, *Am J Physiol Regul Integr Comp Physiol*, 278:R367-R372) demonstrated in animal models that immobilization stress impairs cutaneous barrier recovery and that the glucocorticoid receptor antagonist RU-486 reverses this effect, providing causal evidence that cortisol mediates stress-induced barrier dysfunction. Choi, Brown, Crumrine, and colleagues (2005, *J Invest Dermatol*, 124:587-595) confirmed the finding in human studies, showing that interview-induced and academic-examination stress raised endogenous glucocorticoids and impaired both barrier function and stratum corneum integrity. The specific defects included reduced lamellar body secretion, decreased corneodesmosome density, and impaired epidermal lipid synthesis. Topical application of physiologic lipids normalized the effect.

3.3 Recent human evidence

Pujos, Chamayou-Robert, Parat, Bonnet, and colleagues (2024, *J Cosmet Dermatol*) combined an in vivo arm of 40 women aged 35 to 55 with reconstructed-human-epidermis experiments. The in vivo arm found significant skin-roughness and microrelief differences by chronic-stress status. The reconstructed-epidermis arm demonstrated that cortisol exposure at 2.5 uM reduced filaggrin synthesis by 32% and loricrin by approximately 20%, providing the strongest direct mechanistic evidence to date that cortisol compromises the structural proteins underpinning skin barrier integrity.

3.4 Allostatic load and the hair cortisol biomarker

McEwen's allostatic load model provides the integrative framework. McEwen and Stellar (1993, *Arch Intern Med*) and McEwen (1998, *NEJM*) defined allostatic load as the cumulative wear and tear from repeated activation of stress mediators above their adaptive ranges. The 2010 integrative review with Gianaros extended the model to brain plasticity. Hair cortisol concentration (Stalder 2017 meta-analysis) reflects cumulative cortisol exposure over weeks to months and is the biomarker of choice for chronic-stress intervention validation.

4. THE MULTISENSORY PROTOCOL: ARCHITECTURE AND COMPONENTS

The integrated multisensory protocol examined in this synthesis is delivered in two complementary modalities: an in-person flagship experience and a take-home product extension. The in-person modality engages all five sensory channels through dedicated calibrated components. The take-home extension is anchored in a topical skincare portfolio that itself delivers tactile engagement (through affective self-application), olfactory engagement (through aromatic topical actives), and microbiome restoration (through postbiotic ferments). Figure 4 summarizes the mapping.

Figure 4. The integrated multisensory protocol delivered in two complementary modalities. The in-person component engages all five sensory channels through dedicated calibrated elements. The take-home component is anchored in the topical skincare portfolio, supplemented by an auditory take-home element for habitual engagement.

4.1 Why two integrated modalities

The two modalities are complementary, not redundant. The in-person ritual maximizes intensity, social presence, and sensory richness, three variables associated with the strongest cortisol-lowering effects in the literature. Linnemann, Strahler, and Nater (2016, Psychoneuroendocrinology) demonstrated that the cortisol-reducing effect of music is enhanced when listening occurs in the presence of others. Schofield and Payne (2003) found that immersive multisensory environments outperform quiet rooms for anxiety reduction. The in-person modality functions as the high-intensity dose.

The take-home modality is designed for high frequency. Packheiser and colleagues (2024) identified session frequency as a positive moderator of touch-intervention effectiveness for mental-health outcomes. The take-home extension is anchored in the topical portfolio, which is itself a multi-channel delivery vehicle: it engages affective touch during application, delivers olfactory exposure through aromatic actives, and supplies postbiotic ferments that restore the microbiome substrate disrupted by chronic cortisol. The combined two-modality protocol therefore delivers both clinical-intensity sensory dosing and daily reinforcement frequency.

4.2 Cross-modal component mapping

Table 1. Complete mapping of the integrated multisensory protocol components by sense, modality, and primary literature anchor.

Table 1. provides the complete cross-modal mapping with the primary literature anchor for each component.

SENSE	IN-PERSON COMPONENT	TAKE-HOME COMPONENT	PRIMARY LITERATURE ANCHOR
Auditory	Curated, listener-congruent music played throughout the experience.	Auditory take-home element: hand-wound mechanism producing curated melody, providing combined auditory and haptic engagement.	de Witte 2020, 2022; Linnemann 2015
Tactile	Professional skincare application with facial massage at CT-afferent-optimized stroking velocity.	Daily skincare application: affective self-touch delivering the topical active portfolio.	Packheiser 2024; Loken 2009; Pawling 2017
Gustatory	Hot tea served at the start of treatment as an in-store ritual element.	YouthGlow Collagen Jelly (marine collagen peptides + peach concentrate, daily stick): the nutricosmetic take-home extension supporting hydration, elasticity, and inner-glow via the gut-skin axis.	Pu 2023; de Miranda 2021; Proksch 2014; Bowe and Logan 2011
Olfactory	Calibrated ambient aromatherapy maintained throughout the treatment space.	Aromatic topical actives in serums and creams deliver olfactory engagement during application.	Lehrner 2005; Goel 2005; Sowndhararajan 2016
Visual	Curated visual environment: warm low-illuminance lighting, biophilic elements, sensory congruence.	Intentional self-gaze prompted by the take-home elements; home environment provides extended visual context.	Ulrich 1984; Antonelli 2019; Petrowski 2021

5. EVIDENCE BY SENSORY CHANNEL

This section reviews the peer-reviewed evidence supporting each channel of the integrated protocol. Subsections 5.1 through 5.5 cover the sensory channels of the in-person experience; subsection 5.6 covers the topical, neurocosmetic, and microbiome actives in the take-home portfolio, which together constitute the biochemical channel and the principal vehicle of the take-home extension. Figure 5 provides a master forest plot of pooled effect sizes across the principal modalities; Figures 6 through 9 expand this into per-channel forest plots; Figure 10 plots study quality against effect size to illustrate that larger studies converge on robust moderate-to-large effect magnitudes.

Figure 1. Master forest plot - cortisol, stress, and skin-relevant outcomes across the BONJIL Ritual Channels




Figure 5. Master forest plot of pooled effect sizes for the principal interventions reviewed. Square markers indicate point estimates; horizontal bars denote 95% confidence intervals; marker size is proportional to study or meta-analysis sample size.

Figure 16. Study quality (sample size) vs. effect size across BONJIL Ritual evidence base. Larger studies tend toward more moderate, defensible effects..




Figure 10. Study quality (sample size, log scale) plotted against effect size across the reviewed evidence base. The largest meta-analyses converge on robust moderate effects, with the strongest studies clustering in the medium-to-large effect range.




Figure 32. Bubble chart of the principal meta-analyses cited in this synthesis, plotted by sample size and pooled effect size. Bubble area is proportional to sample size. The cluster of large bubbles in the medium-to-large effect range represents the highest-confidence evidence.

Figure 33. Scatter plot of effect size by publication year for the cited intervention studies. Dot size reflects sample size; the dashed line is the linear trend (slope: +0.015 per year).




Figure 33. Scatter plot of effect size against publication year. The positive trend reflects methodological refinement and improved intervention specification in more recent literature.

Figure 21. Distribution of effect-size estimates across the BONJIL Ritual evidence base. Modal range = 0.3 - 0.6 (small-to-medium); long right tail driven by aromatherapy/sleep




Figure 22. Distribution of participants across the principal sensory-channel meta-analyses synthesized in this review (cumulative $n = 29,522$)




Figure 22. Distribution of participants across the principal sensory-channel meta-analyses synthesized in this review. The cumulative sample totals approximately 29,500 across the five principal modalities.

Figure 22 demonstrates the participant-weighted concentration of the evidence base in the touch and music literatures, which together account for more than three-quarters of pooled participants. The smaller participant counts in the aromatherapy, nature, and sauna literatures should not be interpreted as weaker evidence: each of those literatures contains high-quality randomized controlled trials with large effect sizes that compensate for the smaller cumulative sample. Figure 23 reframes the same evidence base by the mechanism pathways engaged.

Figure 23. Distribution of mechanistic effort across the four principal biological pathways engaged by the multisensory protocol




Figure 23. Distribution of mechanistic effort across the four principal biological pathways engaged by the multisensory protocol. The vagal/parasympathetic pathway accounts for the largest share of the synthesized evidence, followed by central HPA-axis suppression.

5.1 Auditory: structured music and habitual auditory engagement

5.1.1 Mechanism

Auditory input projects from the cochlea through the auditory thalamus to the primary auditory cortex, with collateral projections to limbic and reward structures (amygdala, nucleus accumbens, ventral tegmental area). Music engages dopaminergic reward circuitry: Salimpoor et al. (2011, *Nature Neuroscience*) demonstrated dopamine release in the caudate and nucleus accumbens during music listening. Music simultaneously modulates autonomic balance: relaxing music increases high-frequency heart-rate variability (HF-HRV), a marker of vagal and parasympathetic tone, and reduces salivary alpha-amylase, a marker of sympathetic activation. Chanda and Levitin (2013, *Trends Cogn Sci*) provided the canonical theoretical model linking music to four neurochemical systems: reward, stress and arousal, immunity, and social affiliation.

5.1.2 Primary evidence

de Witte, Spruit, van Hooren, Moonen, and Stams (2020, Health Psychology Review) performed two multilevel meta-analyses on 104 randomized controlled trials of music interventions, involving 9,617 participants and 327 effect sizes. The pooled effect on physiological stress markers, including cortisol, was Cohen's $d = 0.380$; the effect on psychological stress was $d = 0.545$. Among physiological outcomes, heart rate showed the largest effect ($d = 0.456$), followed by blood pressure ($d = 0.343$) and the hormone subset including cortisol ($d = 0.349$).

de Witte and colleagues' subsequent meta-analysis specifically of music therapy (2022, Health Psychology Review, 16:134-159) pooled 47 studies and 2,747 participants, yielding an overall effect size of $d = 0.723$ (95% CI 0.51 to 0.94), a medium-to-large effect. Music therapy outperformed music medicine, with effects larger in non-Western settings and against waiting-list controls.

Linnemann, Ditzen, Strahler, Doerr, and Nater (2015, Psychoneuroendocrinology, 60:82-90) provided ecologically valid demonstration of music's stress-reducing effect. In a daily-life ambulatory study, salivary cortisol and subjective stress were lowest when participants reported listening to music for the purpose of relaxation ($p \leq 0.001$ for both). The finding identifies intentional listening as a key moderator, a moderator that the structured protocol engineers through context, curation, and ritual repetition.

Figure 10. Forest plot - Music interventions across cortisol, autonomic, and psychological outcomes




Figure 6. Forest plot of music interventions across cortisol, autonomic, and psychological outcomes. Structured music therapy with explicit relaxation framing produces consistently medium-to-large effects.

5.1.3 Optimal protocol parameters

The literature converges on the following parameters for maximum cortisol benefit: explicit relaxation intent (Linnemann 2015); listener-congruent selection (de Witte 2020); a slower tempo associated with parasympathetic activation; a minimum continuous duration of approximately 15 to 20 minutes per session; and repeated daily exposure. The structured protocol implements all four moderators.

5.1.4 Music and heart rate variability

Heart rate variability (HRV) provides the most informative non-invasive measurement of parasympathetic engagement during music listening. High-frequency HRV (HF-HRV) reflects vagal tone, while the LF/HF ratio captures sympathovagal balance. Ellis and Thayer (2010) summarized more than two decades of HRV research demonstrating that slow-tempo music with regular rhythmic structure consistently increases HF-HRV and reduces the LF/HF ratio across populations. da Silva and colleagues (2014) demonstrated significant HF-HRV elevation during 30-minute classical-music listening sessions in healthy young adults, with effects sustained for at least 15 minutes post-listening. Trappe (2010, *Heart*, 96:1868-1871) reviewed the cardiovascular literature and concluded that music with tempos near the natural resting heart rate (60 to 80 BPM) produces the strongest entrainment of cardiovascular parameters toward parasympathetic dominance.

Bernardi, Porta, and Sleight (2006, *Heart*, 92:445-452) provided the most elegant demonstration of music's autonomic effects. Across multiple musical genres and tempos, slower tempos and instrumental music produced relaxation responses (reduced ventilation, lower blood pressure, decreased heart rate), while faster tempos and complex rhythms produced opposite effects. Crucially, two-minute pauses between music segments produced even greater relaxation than the music itself, an observation that has informed the design of structured music protocols with built-in rest intervals.

5.1.5 Music and the dopaminergic reward system

The dopaminergic reward system provides a second, complementary mechanism for music's effects on cortisol and wellbeing. Salimpoor, Benovoy, Larcher, Dagher, and Zatorre (2011, *Nature Neuroscience*, 14:257-262) used positron emission tomography with [¹¹C]raclopride to demonstrate distinct dopamine release in the caudate during anticipation of peak emotional moments and in the nucleus accumbens during the peak emotional response itself. The temporal dissociation between anticipatory and consummatory dopamine release parallels the structure of reward processing in other reinforcing stimuli, including food and addictive substances. The implication for ritual design is direct: predictable structure with reliable peak moments produces the largest dopaminergic engagement, supporting the use of curated playlists with established emotional arcs over random or novel music selection.

Koelsch (2014, *Nat Rev Neurosci*, 15:170-180) consolidated the neuroscience of music into a comprehensive framework in which music engages widely distributed networks including auditory cortex, limbic structures (amygdala, hippocampus, hypothalamus), reward and motivation circuits (ventral striatum, ventral tegmental area), and frontal regions (orbitofrontal cortex, ventromedial prefrontal cortex). The breadth of music's neural engagement explains its capacity to influence cortisol through multiple parallel mechanisms: direct autonomic modulation via brainstem and limbic projections, reward-system engagement via mesolimbic dopamine, and top-down cognitive regulation via prefrontal cortex.

5.1.6 Clinical applications in surgical and medical settings

Music therapy has been validated in multiple clinical contexts directly relevant to cortisol-reduction protocol design. Bradt and Dileo's Cochrane reviews of music for coronary heart disease (2009, updated 2013) and music for mechanically ventilated patients (2014) documented consistent reductions in anxiety, heart rate, blood pressure, and respiratory rate. Pelletier (2004, *J Music Ther*, 41:192-214) performed an early meta-analysis demonstrating that music alone and music combined with relaxation techniques both significantly decreased arousal and stress.

The integration into structured multisensory protocols draws on this clinical foundation: music functions as both an autonomic regulator (heart rate variability and blood pressure) and a cognitive primer (attention, expectation, and mood). When deployed in the context of a deliberate sensory environment with paired olfactory, tactile, and visual cues, music's effects are amplified through the convergent engagement of multiple limbic and brainstem targets.

5.2 Tactile: skincare with massage and affective self-application

5.2.1 Mechanism


Gentle stroking of hairy skin activates C-tactile (CT) afferents, unmyelinated low-threshold mechanoreceptors with conduction velocities of approximately 0.6 to 1.3 m/s that project preferentially to the posterior insular cortex, the dedicated affective-touch pathway. CT afferents demonstrate an inverted-U response curve to stroking velocity, with peak firing and peak subjective pleasantness at approximately 3 cm/s and broader optimal activation across the 1 to 10 cm/s range (Loken, Wessberg, Morrison, McGlone, and Olausson, 2009, *Nature Neuroscience*, 12:547-548). Vallbo, Olausson, and Wessberg (1999, *J Neurophysiol*, 81:2753-2763) established CT afferents as a second tactile system in hairy skin, with mechanical thresholds as low as 0.04 mN. McGlone, Wessberg, and Olausson (2014, *Neuron*, 82:737-755) consolidated the framework distinguishing discriminative (myelinated A-beta) from affective (unmyelinated CT) touch.

Touch additionally engages the oxytocinergic system. Uvnas-Moberg, Handlin, and Petersson (2015, *Frontiers in Psychology*, 5:1529) reviewed the neuroendocrine cascade: touch leads to oxytocin release, which inhibits CRF and ACTH, which reduces adrenal cortisol output. The system is bidirectional: touch directly engages parasympathetic vagal tone while damping the HPA axis upstream.

5.2.2 Primary evidence: the Packheiser 2024 meta-analysis

Packheiser, Hartmann, Fredriksen, Gazzola, Keyzers, and Michon (2024, *Nature Human Behaviour*, 8:1088-1107) performed a pre-registered systematic review (PROSPERO CRD42022304281) and multivariate meta-analysis of 137 touch-intervention studies encompassing 12,966 participants across 166 cohorts and 643 effect sizes. The overall effect of touch was Hedges' $g = 0.52$ (95% CI 0.42 to 0.63). Channel-specific effects are presented in Figure 7 and include cortisol regulation in newborns ($g = 0.78$, 95% CI 0.24 to 1.31), weight gain in newborns (0.65, 0.37 to 0.94), pain reduction in adults (0.69, 0.48 to 0.89), depression (0.59, 0.40 to 0.78), state anxiety (0.64, 0.44 to 0.84), and trait anxiety (0.59, 0.40 to 0.77).

Packheiser's analysis identified several moderators directly relevant to multisensory protocol design: human touch outperformed object-mediated and robot touch for mental-health outcomes; session frequency was a positive moderator for mental-health outcomes; and head and face touch produced larger benefits than touch to other body regions. The implications for facial skincare protocol design are direct: frequent, deliberate, head and face affective self-application is positioned as the highest-leverage tactile intervention available.

Figure 9. Forest plot - Touch interventions across cortisol, anxiety, depression, pain, and weight gain**Figure 7.** Forest plot of touch interventions across cortisol, anxiety, depression, pain, and weight-gain outcomes. The Packheiser (2024) sub-analyses dominate, supplemented by experimental work from Ditzen (2007), Holt-Lunstad (2008), and Pawling (2017).

5.2.3 Additional supporting studies

Ditzen and colleagues (2007, *Psychoneuroendocrinology*, 32:565-574) provided RCT-level evidence for the specific cortisol benefit of pre-stressor neck and shoulder massage: salivary cortisol response to a subsequent laboratory stressor was significantly attenuated relative to no-touch and verbal-support control conditions. Holt-Lunstad, Birmingham, and Light (2008, *Psychosomatic Medicine*, 70:976-985) extended this with a controlled warm-touch intervention: increased salivary oxytocin, reduced alpha-amylase, and reduced 24-hour ambulatory systolic blood pressure.

5.2.4 CT-afferent stroking parameters

Pawling, Cannon, McGlone, and Walker (2017, *PLoS ONE*, 12:e0173457) confirmed that stroking at 3 cm/s produces greater subjective pleasantness, greater heart-rate deceleration, and selectively greater zygomaticus (smile-muscle) EMG activity than stroking at 30 cm/s on the forearm. Ackerley and colleagues (2014) established that the CT-optimal stimulus is presented at skin temperature; warmer or cooler stroking attenuates the affective response. Croy, Bierling, Sailer, and Ackerley (2021, *Neuroscience*, 464:33-43) characterized individual variability in pleasantness ratings to stroking touch over different velocities. Taneja and colleagues' (2021) systematic review and meta-analysis on defining pleasant touch stimuli confirmed the central role of the 1 to 10 cm/s range.

These data permit a precise specification of optimal skincare-application velocity: deliberate, slow, approximately 3 cm/s strokes across the face during application, performed on hairy (non-glabrous) skin at skin temperature.

5.2.5 The oxytocin-cortisol axis

The mechanistic case for tactile intervention rests heavily on the oxytocin-cortisol relationship. Oxytocin is released in response to non-noxious sensory stimulation, with touch the most reliable trigger in human populations. Uvnas-Moberg, Handlin, and Petersson (2015, *Frontiers in Psychology*, 5:1529) consolidated three decades of research demonstrating that touch-induced oxytocin release produces measurable reductions in CRF and ACTH at the hypothalamic and pituitary levels, with downstream cortisol suppression at the adrenal cortex. Light, Grewen, and Amico (2005, *Biol Psychol*, 69:5-21) demonstrated that women who reported frequent partner hugs showed both higher baseline oxytocin and lower blood pressure than women with less frequent partner contact, providing population-level evidence for the touch-oxytocin-cardiovascular cascade.

Heinrichs, Baumgartner, Kirschbaum, and Ehlert (2003, *Biol Psychiatry*, 54:1389-1398) provided experimental confirmation: intranasal oxytocin administration combined with social support produced the largest reductions in salivary cortisol response to the Trier Social Stress Test, larger than either oxytocin alone or social support alone. The finding indicates that oxytocin and social-contextual stress reduction work synergistically rather than additively, with implications for protocol design: deliberate engagement of touch in a supportive social context produces effects larger than the sum of the individual components.

5.2.6 Field and the Touch Research Institute

The Touch Research Institute (Field T, Miller School of Medicine, University of Miami) has produced more than two hundred studies on touch and massage across populations including preterm infants, adults with chronic pain, individuals with depression, and patients with cancer. Field, Diego, and Hernandez-Reif (2007, *Infant Behav Dev*, 30:367-383) reviewed the preterm-infant literature, documenting that moderate-pressure massage produced significant weight gain (approximately 47% greater than control infants in the largest studies), shorter hospital stays, and improved developmental outcomes at follow-up. The biological mechanism, as detailed in Field's subsequent reviews, involves increased vagal activity, increased gastric motility, and reduced cortisol production.

Field, Hernandez-Reif, Diego, Schanberg, and Kuhn (2005, *Int J Neurosci*, 115:1397-1413) performed a quantitative review of cortisol and serotonin changes following massage therapy across multiple populations and clinical conditions. The pooled analysis demonstrated an average 31% reduction in cortisol and average 28 to 31% increases in serotonin and dopamine following massage therapy interventions, providing benchmark effect sizes that have informed subsequent protocol design.

5.3 Gustatory: the warm-beverage ritual and the nutricosmetic take-home extension

5.3.1 In-person component: the warm-beverage ritual

The in-person component delivers the gustatory channel as a hot tea served at the start of treatment. The contribution is principally non-pharmacological. The act of slow, warm, intentional consumption activates parasympathetic vagal tone through several converging pathways: warm visceral input from gastric receptors, deliberate breath pacing during drinking, and the ritual-cognitive framing that prepares the participant for the treatment that follows. To the extent that the tea selected is a *Camellia sinensis* preparation, the L-theanine content provides a modest acute parasympathetic effect (White et al., 2016, *Nutrients*, 8:53). The dominant mechanism, however, is ritual psychology: the warm-beverage ritual serves as a conditioned cue establishing the transition from external-world stress to the protocol's parasympathetic-engagement state.

5.3.2 Take-home component: *BONJIL YouthGlow Collagen Jelly*

The take-home extension of the gustatory channel is delivered as BONJIL YouthGlow Collagen Jelly, a daily marine collagen peptide and peach concentrate jelly stick. The take-home ingestible channel is positioned to perform three complementary functions: (a) sustain the daily-ritual habit established by the in-person tea ceremony, (b) deliver bioactive nutricosmetic support for the dermal substrate damaged by chronic cortisol exposure, and (c) engage the gut-skin axis as a parallel route to the cutaneous outcomes targeted by the broader protocol.

5.3.3 Mechanism: *bioactive collagen peptides from gut to skin*

Hydrolyzed marine collagen is absorbed as bioactive di-peptides and tri-peptides rather than as intact protein. Iwai and colleagues (2005, *J Agric Food Chem*, 53:6531-6536) provided the foundational human bioavailability evidence: after ingestion of 9.4 to 23 g gelatin hydrolysate, hydroxyproline-containing peptides rose to a maximum of 20 to 60 nmol/mL plasma at 1 to 2 hours, with proline-hydroxyproline (Pro-Hyp) constituting approximately 50% of food-derived collagen peptides detected in blood. The imino-acid ring structures of proline and hydroxyproline resist brush-border and cytoplasmic peptidases, allowing the bioactive peptides to survive digestion intact.

Yazaki and colleagues (2017, *J Agric Food Chem*, 65:2315-2322) closed the bioavailability loop by demonstrating transport of glycyl-prolyl-hydroxyproline (Gly-Pro-Hyp) and Pro-Hyp into the bloodstream and the skin, identifying approximately seventeen collagen-derived peptides with enrichment in skin tissue. Ohara and colleagues (2010, *J Dermatol*, 37:330-338) established the dermal mechanism: Pro-Hyp at physiologically achievable concentrations enhanced human dermal fibroblast proliferation 1.5-fold and hyaluronic acid synthesis 3.8-fold, with a 2.3-fold elevation of HAS2 messenger RNA. Small-interfering-RNA knockdown of HAS2 abolished both the HAS2 transcription effect and the mitotic effect, confirming a signaling-mediated mechanism rather than substrate provision.

Proksch and colleagues (2014, *Skin Pharmacol Physiol*, 27:113-119) confirmed the downstream histological effect in suction-blister biopsies from a randomized controlled trial of 114 women: oral collagen peptides at 2.5 g per day for 8 weeks produced a 65% increase in type I procollagen and an 18% increase in elastin in the dermal extracellular matrix versus placebo. The combined mechanistic chain, ingestion to plasma to skin to fibroblast to procollagen and hyaluronic acid synthesis, is one of the most completely documented bioactive food-derived ingredient pathways in current dermatological nutrition science.

5.3.4 Primary clinical evidence: *meta-analytic consensus*

Three independent meta-analyses provide the headline clinical case. Pu, Huang, Pu, Kang, Hoang, Chen, and Chen (2023, *Nutrients*, 15:2080) pooled 26 randomized controlled trials with 1,721 participants. The hydration effect was significant ($Z = 4.94$, $p < 0.00001$). The elasticity effect across 19 studies was a pooled effect size of 0.72 (95% CI 0.40 to 1.03; $Z = 4.49$, $p < 0.00001$), placing collagen peptide supplementation in the medium-to-large effect range for elasticity outcomes.

de Miranda, Weimer, and Rossi (2021, *Int J Dermatol*, 60:1449-1461) pooled 19 randomized, double-blind, controlled trials with 1,125 participants, ages 20 to 70, 95% women. Grouped analysis favored hydrolyzed collagen over placebo for hydration, elasticity, and wrinkles, with hydration and elasticity confirmed in subgroup meta-analysis. Effective daily doses spanned 2.5 to 10 g, with the conclusion that ingestion of hydrolyzed collagen for 90 days reduces skin aging. Danessa, Notario, and Regina (2025, *Indian J Dermatol Venereol Leprol*, 91:730-740) confirmed the pattern in a more selective meta-analysis: hydration SMD 1.25 (95% CI 0.77 to 1.74) and elasticity SMD 0.61 (95% CI 0.21 to 1.02). Most common dose was 4 g per day; effects emerged consistently after 8 weeks.

5.3.5 Marine-specific evidence

Marine collagen is the most studied source for nutricosmetic applications and is associated with the highest amino-acid homology to human type I collagen, the dominant dermal collagen. Kim, Chung, Choi, Sakai, and Lee (2018, *Nutrients*, 10:826) demonstrated in a randomized, double-blind, placebo-controlled trial of 64 participants that a fish-derived low-molecular-weight collagen peptide at 1,000 mg per day for 12 weeks significantly improved hydration at 6 and 12 weeks, improved elasticity, and significantly improved three wrinkle parameters at 12 weeks, with no adverse effects.

Inoue, Sugihara, and Wang (2016, *J Sci Food Agric*, 96:4077-4081) demonstrated improved facial moisture, improved elasticity, and reduced facial aging signs with fish collagen hydrolysate in a randomized double-blind placebo-controlled study. Koizumi, Inoue, Shimizu, and colleagues (2018, *Int J Pept Res Ther*, 24:397-402) demonstrated that 3,000 mg per day of fish-scale-derived collagen peptides for 12 weeks in 71 women aged 30 to 60 significantly decreased periorbital wrinkles ($p < 0.05$) and enhanced facial moisture and elasticity ($p < 0.001$ for both) with no adverse events. Asserin, Lati, Shioya, and Prawitt (2015, *J Cosmet Dermatol*, 14:291-301) demonstrated significantly increased dermal collagen density and significantly decreased dermal collagen network fragmentation by 4 weeks of supplementation at 10 g per day, persisting to 12 weeks, the first clinical evidence of an effect on dermal collagen fragmentation. Lin and colleagues (2021, *J Cosmet Dermatol*, 20:188-194) demonstrated, in a randomized placebo-controlled trial of 50 volunteers over 8 weeks, hydration +17.8%, brightness +5.4%, crow's feet +14.9%, texture +9.9%, wrinkles +29.3%, and collagen content +22.3% from baseline.

Maia Campos, Franco, Kakuda, Cadioli, Costa, and Bouvret (2021, *Molecules*, 26:4880) demonstrated in a double-blind placebo-controlled clinical study that hydrolyzed fish cartilage containing greater than 65% low-molecular-weight collagen peptides at 500 mg per day for 90 days significantly reduced wrinkles, significantly increased dermal echogenicity and thickness, and improved collagen morphology and reduced elastosis on reflectance confocal microscopy.

5.3.6 Peach concentrate: the vitamin C and antioxidant adjunct

Peach (*Prunus persica*) supplies the gustatory channel with two complementary biological functions beyond its role as a palatable carrier. First, peach contains 4 to 13 mg of vitamin C per 100 g (Gil, Tomas-Barberan, Hess-Pierce, and Kader, 2002, *J Agric Food Chem*, 50:4976-4982). Vitamin C is an essential cofactor for prolyl and lysyl hydroxylase, the enzymes that catalyze the post-translational hydroxylation of proline and lysine residues required for stable triple-helix collagen formation (Murad, Grove, Lindberg, Reynolds, Sivarajah, and Pinnell, 1981, *Proc Natl Acad Sci USA*, 78:2879-2882). The pairing of marine collagen peptides (substrate and signaling) with peach-derived vitamin C (cofactor) is therefore directly aligned with the collagen-biosynthetic pathway.

Second, peach supplies a documented carotenoid and polyphenol antioxidant profile: total carotenoids 71 to 210 ug per 100 g, dominated by beta-carotene (approximately 88% of total carotenoids; Belisario et al., 2023, *PMC10039266*) and beta-cryptoxanthin, with phenolic compounds dominated by chlorogenic acid and (+)-catechin (Aubert and Chalot, 2020, *J Food Compos Anal*, 86:103337). Dietary carotenoids distribute into light-exposed skin and provide systemic antioxidant photoprotection against ultraviolet-mediated erythema with sustained intake of approximately 10 weeks or longer (Stahl and Sies, 2012, *Am J Clin Nutr*, 96:1179S-1184S). The peach concentrate functions as a continuous, gentle systemic antioxidant load supporting the broader protocol's stress-reduction architecture.

5.3.7 The gut-skin axis bridge

The ingestible channel is uniquely positioned to engage the gut-skin axis in parallel with the topical and sensory channels of the broader protocol. Bowe and Logan (2011, *Gut Pathogens*, 3:1) demonstrated that the gut microbiome regulates systemic inflammation, oxidative stress, and skin homeostasis, with probiotic supplementation associated with lower cortisol and reduced anxiety in healthy adults. The combination of bioactive marine peptides (supporting fibroblast function), vitamin C (supporting collagen biosynthesis), and carotenoid and polyphenol antioxidants (supporting systemic redox balance) delivers a coordinated nutricosmetic intervention that complements the sensory and topical channels of the multisensory protocol at the systemic, from-within level. The result is a true five-channel convergent intervention: the gustatory channel becomes a daily nutritional reinforcement of the same dermal substrate and stress-modulation pathways engaged in the in-person experience.

5.3.8 Dose, format, and timing

The literature converges on a daily dose of approximately 2.5 to 10 g of hydrolyzed marine collagen peptides, with low-molecular-weight peptide preparations effective at 1 g per day. Time to effect is approximately 4 to 8 weeks for hydration outcomes, 8 to 12 weeks for elasticity and dermal density, and 12 weeks or longer for visible wrinkle outcomes. Benefits partly persist after discontinuation (Proksch 2014; Sangsuwan 2021; Bolke 2019), supporting a sustained daily maintenance model rather than intermittent or loading-dose use. The jelly stick format selected for YouthGlow optimizes palatability, portability, and adherence to the daily ritual cadence that the multisensory protocol requires for sustained effect.

5.4 Olfactory: aromatherapy and topical olfactory delivery

5.4.1 Mechanism: direct limbic access

Olfactory input is anatomically unique among the sensory channels. The olfactory bulb projects directly to the piriform cortex, amygdala, hippocampus, and entorhinal cortex without the thalamic relay required by all other sensory modalities. This direct limbic access explains the exceptional speed and emotional intensity of olfactory experience and the well-documented Proust effect, in which odors trigger more vivid and emotionally laden autobiographical memories than visual or verbal cues. Larsson, Willander, Karlsson, and Arshamian (2014, *Front Psychol*, 5:1011) reviewed the olfactory memory literature and confirmed that odor-evoked memories are characterized by greater emotional intensity and earlier life reference points than memories evoked through other modalities. Saive, Royet, and Plailly (2014, *Front Behav Neurosci*, 8:240) extended this work to demonstrate that olfactory-triggered autobiographical retrieval engages limbic networks more strongly than non-olfactory triggers, with implications for ritual design: scent-anchored ritual experiences establish stronger emotional and memory traces than non-scented equivalents.

Soudry, Lemogne, Malinvaud, Consoli, and Bonfils (2011, *Eur Ann Otorhinolaryngol Head Neck Dis*, 128:18-23) detailed the anatomical projections in clinically oriented terms: olfactory input reaches the orbitofrontal cortex for conscious identification, the amygdala for emotional valuation, and the hippocampus for memory encoding, all within approximately 200 to 400 milliseconds of stimulus presentation. The combination of direct limbic access, autonomic engagement, and memory encoding makes olfaction the most efficient sensory channel for establishing conditioned associations between a stimulus and a desired physiological state.

5.4.2 *Silexan: the strongest oral aromatherapy evidence*

Silexan, an orally administered preparation of *Lavandula angustifolia* essential oil standardized to specific terpene composition, has accumulated the strongest evidence base for olfactory-pathway anxiolysis. Yap, Dolzhenko, Jalal, Hadi, and Khan (2019, *Scientific Reports*, 9:18042) performed a network meta-analysis of five randomized controlled trials with 524 participants on Silexan 80 mg and 121 on 160 mg. Silexan 160 mg outperformed 80 mg, placebo, and paroxetine on Hamilton Anxiety Rating Scale (HAMA) reduction, with surface under the cumulative ranking curve (SUCRA) values placing Silexan 160 mg as the best-ranked intervention across the network.

Dold, Bartova, Volz, Seifritz, Moller, Schlafke, and Kasper (2023, *Eur Arch Psychiatry Clin Neurosci*, 273:1615-1628) extended the evidence base with a meta-analysis of randomized placebo-controlled trials confirming significant HAMA reductions and identifying linalool and linalyl acetate as the principal active terpenes. Kasper, Gastpar, Muller, Volz, Moller, Dienel, and Schlafke (2014, *Int Clin Psychopharmacol*, 29:269-277) demonstrated that Silexan was non-inferior to lorazepam in subsyndromal anxiety, an important comparative-effectiveness finding given lorazepam's established efficacy.

The Silexan mechanism, as elucidated through preclinical work by Schuwald and colleagues (2013) and confirmed in human pharmacodynamic studies, involves modulation of voltage-gated calcium channels in serotonergic, noradrenergic, and glutamatergic neurons. The terpene components also interact with the GABA system, though through allosteric modulation rather than direct GABA-A receptor binding, explaining the absence of sedation and dependence potential characteristic of benzodiazepines.

5.4.3 *Inhalation aromatherapy and sleep*

Hwang and Shin (2015, *J Altern Complement Med*, 21:61-68) meta-analyzed twelve randomized controlled trials of aromatherapy on sleep quality. Pooled standardized mean difference was 1.14 (95% CI 0.54 to 1.74; $Z = 3.716$), a large effect size. Subgroup analysis demonstrated that inhalation methods (95% CI 0.79 to 1.54; $Z = 6.107$) outperformed aromatherapy massage (95% CI 0.13 to 2.17; $Z = 2.205$), an important practical finding given that inhalation is the simpler and more readily implemented modality.

Lillehei and Halcon (2014, *J Altern Complement Med*, 20:441-451) performed an independent systematic review of essential oil inhalation for sleep, reaching consistent conclusions: lavender essential oil produced the most consistent improvements across sleep outcomes, with smaller but significant effects for chamomile, ylang ylang, and bergamot preparations. The literature supports the use of inhaled aromatherapy as both an acute sleep-onset aid and a chronic sleep-quality enhancer when applied consistently over several weeks.

5.4.4 *Bergamot and acute parasympathetic activation*

Watanabe, Kuchta, Kimura, Rauwald, Kamei, and Imanishi (2015, *Forschende Komplementarmedizin*, 22:43-49) demonstrated that 15 minutes of bergamot essential oil inhalation increased high-frequency heart rate variability and reduced salivary cortisol in healthy adults, with effects emerging within the first 5 minutes and sustained throughout the exposure period. The bergamot preparation, derived from *Citrus bergamia* peel and containing limonene, linalyl acetate, linalool, and bergapten, produces parasympathetic activation through dual mechanisms: direct GABA modulation by the linalool component and olfactory-pathway limbic engagement.

Saiyudthong and Marsden (2011) demonstrated bergamot oil's anxiolytic effects in animal models, and Han, Cho, and Cho (2017) extended this work to human populations, demonstrating salivary alpha-amylase reductions consistent with sympathetic-nervous-system attenuation. The convergent evidence supports bergamot as an effective acute parasympathetic activator with rapid onset, ideal for the opening phase of structured multisensory protocols.

5.4.5 Vanilla and conditioned safety responses

Lehrner, Marwinski, Lehr, Jöhren, and Deecke (2005, *Physiol Behav*, 86:92-95) demonstrated that ambient vanilla and orange odors in a dental waiting room significantly reduced anxiety in female patients (effect size $d = 0.42$), with smaller but significant effects in males. The vanilla finding is mechanistically interesting because vanilla is one of the earliest and most universally encountered odors in human development, present in breast milk and reliably associated with feeding and maternal contact in infancy. Schaal and colleagues (2000) demonstrated that newborn infants preferentially orient to vanilla odor within the first hours of life, indicating a hardwired conditioning between vanilla and safe feeding states.

The use of *Vanilla Planifolia* Leaf Cell Extract in topical formulations provides a double-pathway intervention: the olfactory exposure during application activates the vanilla-safety associations established in infancy, while the bioactive constituents (vanillin, vanillic acid, and polyphenols) provide direct antioxidant activity at the skin surface. This dual function exemplifies the integrated design philosophy underlying the multisensory protocol: each component contributes through multiple parallel mechanisms.

5.4.6 Lavender, salivary cortisol, and cortisol-awakening response


Toda and Morimoto (2008, *Arch Oral Biol*, 53:964-968) demonstrated that lavender aroma exposure significantly reduced salivary cortisol in healthy adults within 10 minutes of exposure, with effects largest in subjects with higher baseline cortisol values. The preferential effect on high-cortisol individuals is consistent with the broader literature on stress-reduction interventions: subjects with the greatest physiological stress burden show the largest absolute benefits from intervention.

Sayorwan, Siripornpanich, Piriapunyaporn, Hongratanaworakit, Kotchabhakdi, and Ruangrunsi (2012, *J Med Assoc Thai*, 95:598-606) extended this work by simultaneously measuring EEG, autonomic activity, and mood after lavender inhalation. The combined measurement demonstrated increased theta power in the frontal regions (associated with reduced cognitive load and increased relaxation), reduced systolic blood pressure, and improved self-reported calmness, indicating that lavender's effects span cortical, autonomic, and subjective levels simultaneously.

5.4.7 The Sowndhararajan synthesis

Sowndhararajan and Kim (2016, *Sci Pharm*, 84:724-751) provided the most comprehensive available review of fragrance influences on human psychophysiology. The review synthesized research across more than fifty essential oils and fragrance compounds, identifying lavender, rose, bergamot, ylang-ylang, sandalwood, and frankincense as the most consistently anxiolytic, with smaller bodies of evidence supporting jasmine, neroli, and rosemary. Across the reviewed literature, the consistent pattern was rapid onset (within 5 to 15 minutes of exposure), modest-to-large effect sizes, and dose-response relationships in which low-to-moderate olfactory loads outperformed high-intensity exposures.

The dose-response finding has direct implications for protocol design: calibrated, low-to-moderate ambient aromatic loads sustained throughout an experience produce larger effects than brief high-intensity exposures. This informs the design of the in-person treatment-space aromatic environment, which is calibrated to a subliminal-to-comfortable intensity level rather than to maximum perceptible concentration.

Figure 11. Forest plot - Olfactory and aromatherapy interventions across cortisol, sleep, and anxiety**Figure 8.** Forest plot of olfactory and aromatherapy interventions across cortisol, sleep, and anxiety outcomes. Aromatherapy produces consistently moderate-to-large effects, with inhalation outperforming aromatherapy massage and Silexan 160 mg matching standard pharmacotherapy.

5.5 Visual: lighting, biophilic design, and the curated environment

5.1.1 Mechanism

Ulrich's 1984 paper in *Science* remains the founding document of evidence-based biophilic design. The study examined postoperative recovery in patients undergoing cholecystectomy at a Pennsylvania hospital. Patients assigned to rooms with windows facing a stand of deciduous trees had shorter postoperative hospital stays (mean 7.96 days vs 8.70 days, $p = 0.027$), received fewer negative nursing notations regarding their condition, and required less potent analgesic medication than patients assigned to rooms with windows facing a brick wall. The study controlled for surgical technique, age, sex, smoking history, weight, and prior hospitalization. The finding was foundational because it demonstrated, in a quasi-experimental clinical context with hard endpoints, that visual environment produces measurable health outcomes.

Subsequent replications and extensions have confirmed and refined the original finding. Beauchemin and Hays (1996) demonstrated reduced mortality and length of stay in cardiac intensive-care patients in sunny rooms. Walch and colleagues (2005, *Psychosom Med*, 67:156-163) showed that spinal-surgery patients in bright rooms used 22% less analgesic medication and reported lower stress than patients in dim rooms. Park and Mattson (2009, *HortTechnology*, 19:563-568) extended the finding to plants in hospital rooms: appendectomy patients with plants in their rooms required less postoperative pain medication and reported lower anxiety and higher satisfaction than patients without plants.

5.5.2 *Shinrin-yoku and the Park-Miyazaki research program*

The Japanese shinrin-yoku (forest-bathing) research program led by Park, Tsunetsugu, Kasetani, Kagawa, and Miyazaki has produced the most extensive body of field research on biophilic exposure and physiology. Park and colleagues (2010, *Environ Health Prev Med*, 15:18-26) reported the principal field experiment: 280 subjects across 24 forests participated in paired-day experiments comparing forest exposure to urban exposure. Forest exposure produced lower salivary cortisol, lower pulse rate, lower blood pressure, greater parasympathetic nervous activity (HF-HRV), and lower sympathetic nervous activity (LF/HF ratio) than urban exposure. The cortisol reduction averaged approximately 12.4% from pre- to post-exposure in the forest condition, compared to 1.6% in the urban condition.

Antonelli, Barbieri, and Donelli (2019, *Int J Biometeorol*, 63:1117-1134) performed the definitive meta-analysis: 22 studies in systematic review, 8 in the pooled quantitative synthesis. Post-intervention salivary cortisol mean difference was -0.05 ug/dL (95% CI -0.06 to -0.04; $p < 0.01$). Pre-intervention mean difference was -0.08 ug/dL (95% CI -0.11 to -0.05). The consistent and substantial post-exposure cortisol reduction places forest bathing among the most reliably effective single-modality cortisol-reduction interventions.

Li and colleagues' work at Nippon Medical School has extended the field-research base into the immunological domain. Li (2010, *Environ Health Prev Med*, 15:9-17) demonstrated that forest-bathing trips significantly increased natural-killer-cell activity and the intracellular expression of anti-cancer proteins (perforin, granulysin, granzymes), with effects sustained for more than 30 days post-exposure. The proposed mechanism involves phytoncides, the volatile organic compounds released by trees, which have been demonstrated in vitro to enhance NK cell function. The immunological extension reframes forest bathing as not only a cortisol-reduction intervention but a broader systemic physiological optimizer.

5.5.3 *Attention restoration and fractal aesthetics*

Kaplan and Kaplan's Attention Restoration Theory, articulated in *The Experience of Nature* (1989), provides the theoretical framework underpinning the biophilic-design evidence. The framework distinguishes directed attention (effortful, fatigable) from involuntary attention (effortless, restorative). Natural environments, with their inherent richness and gentle stimulation, engage involuntary attention while allowing directed attention to recover. Berman, Jonides, and Kaplan (2008, *Psychol Sci*, 19:1207-1212) provided experimental confirmation: 50-minute walks in nature produced significantly greater improvements in directed-attention tasks than equivalent walks in urban environments.

The aesthetic mechanism underlying nature's restorative effects has been illuminated through fractal-dimension research. Hagerhall, Purcell, and Taylor (2004, *J Environ Psychol*, 24:247-255) demonstrated that natural scenes with fractal dimensions between approximately 1.3 and 1.5 (the range characterizing most natural landscapes) elicit greater aesthetic preference than scenes with higher or lower fractal complexity. Taylor (2006, *Leonardo*, 39:245-251) extended this work to EEG measurement, demonstrating that viewing 1.3-1.5 fractal images produced increased alpha-band activity (associated with relaxed wakefulness) and reduced stress response compared to viewing higher-dimension fractals. The convergence of aesthetic preference and physiological relaxation around the natural fractal dimension provides a quantitative principle for visual environmental design.

5.5.4 Light spectrum, illuminance, and cortisol

Beyond the content of the visual environment, the physical properties of lighting itself modulate cortisol and autonomic function. Petrowski, Schmalbach, and Strahler (2021, *Stress*, 24:60-69) demonstrated that warm-temperature, dim lighting produced lower post-stress cortisol than bright, cool lighting in a controlled experimental paradigm. The effect was consistent with melanopsin-pathway physiology: bright, blue-enriched light (5000K to 6500K) suppresses melatonin and increases alertness through retinal ganglion cell projections to the suprachiasmatic nucleus and the locus coeruleus, while warm, dim light (2700K to 3000K) supports parasympathetic engagement and the natural evening transition to rest-state physiology.

Smolders, de Kort, and Cluitmans (2012, *Physiol Behav*, 107:7-16) demonstrated dose-dependent effects of illuminance on cortisol awakening response: higher daytime light exposure produced larger cortisol awakening responses (a marker of healthy HPA axis function) the following morning. The combined evidence indicates that visual environment should be designed with explicit attention to both content (natural elements, fractal complexity, biophilic features) and lighting properties (color temperature, illuminance, dynamic variation), with warm low-illuminance environments selected for parasympathetic-engagement contexts and brighter cooler environments reserved for daytime activation contexts.

5.5.5 Biophilic design as a clinical-grade intervention

Browning, Ryan, and Clancy's framework 14 Patterns of Biophilic Design (Terrapin Bright Green, 2014) consolidates the empirical literature into actionable design specifications across three categories: nature in the space (visual connection with nature, non-visual connection with nature, non-rhythmic sensory stimuli, thermal and airflow variability, presence of water, dynamic and diffuse light, connection with natural systems), natural analogues (biomorphic forms and patterns, material connection with nature, complexity and order), and nature of the space (prospect, refuge, mystery, risk/peril). Each pattern is supported by published empirical research with documented cortisol, autonomic, cognitive, or affective benefits.

The integrated multisensory protocol's in-person environment implements multiple biophilic patterns simultaneously: visual connection with natural materials, dynamic and diffuse lighting through warm low-illuminance fixtures, biomorphic forms in furnishings and architecture, the prospect-and-refuge balance characteristic of well-designed spaces, and thermal variability through the integration of warm tea, warm towels, and skin-temperature topical application. The cumulative visual-environmental load is therefore not a single intervention but a layered biophilic design that engages multiple validated stress-reduction patterns simultaneously.

Figure 12. Forest plot - Visual environment and nature exposure on cortisol/autonomic outcomes




Figure 9. Forest plot of visual environment and nature exposure on cortisol and autonomic outcomes.

5.6 The topical active portfolio: convergent mechanism pathways

The topical active portfolio of the take-home component is organized into three convergent mechanism pathways: barrier repair and antioxidant defense; direct neurocosmetic signaling at cutaneous nerve endings and neuromuscular junctions; and microbiome and neuroimmune restoration. Figure 12 maps each active onto its mechanism pathways. Figure 13 presents the portfolio organization.

Figure 15. Ingredient x mechanism heat map for the BONJIL active portfolio.




Figure 12. Ingredient by mechanism heat map for the topical active portfolio. Strength of evidence is graded from blank (no relevant evidence) to ★★★★★ (strong evidence).

BONJIL ACTIVE PORTFOLIO - Three Convergent Mechanism Pathways

BARRIER REPAIR	NEUROCOSMETIC	MICROBIOME & NEUROIMMUNE
Counters Chol 2005 / ujos 2024 cortisol barrier damage	Cutaneous nerve endings + neuromuscular modulation	Postbiotic balance, brain-skin axis restoration
Niacinamide (B3) Bissett 2005, Ilhani 2005	MUNAPSYS [Tyr C] Lipotec Synth Hex-1	Lactobacillus Ferment Komatsu 2005, Roth 2018
Ceramide NP Masukawa 2014	Palmitoyl Tripeptide-1 Patent 2008/2018	Beta-Glucan Du 2014
Squalane Emollient lipid layer	Tetrapeptide-7 Pentaplex 2008	Biosaccharide Gum-1 Humectant, little efficacy data
Panthenol (B5) Ehlers 2003	Ectoin March 2014, Buenger 2007	BiomeGlow Blue Postbiotic protocol
Sodium Hyaluronate Humectant	Vanilla Planifolia [Tyr C] Lebreton 2005 (scent)	Turmeric Extract Vaughn 2016
Vitamin E / Vit C Oxidative stress	Helichrysum [Tyr C] SILAB in-house only	Calendula Extract Della Loggia, Fiumi
Convergent cortisol-axis attenuation + barrier-substrate restoration		

Figure 13. The topical portfolio organized into three convergent mechanism pathways. The portfolio is designed for substrate restoration of barrier structures damaged by chronic cortisol, neurocosmetic signaling at cutaneous nerve endings, and microbiome rebalance.

5.6.1 Barrier repair and antioxidant actives

Niacinamide (vitamin B3). Bissett, Miyamoto, Sun, Li, and Berge (2004, Int J Cosmet Sci, 26:231-238) performed a 12-week double-blind, placebo-controlled, split-face RCT in 50 women aged 40 to 60. Topical 5% niacinamide significantly reduced yellowing, wrinkling, red blotchiness, and hyperpigmented spots. Bissett, Oblong, and Berge's tollow-up (2005, Dermatol Surg, 31:860-865) consolidated the dose-response data summarized in Figure 14. Hakozaiki and colleagues (2002, Br J Dermatol, 147:20-31) demonstrated the mechanism for the pigmentation benefit: suppression of melanosome transfer from melanocytes to keratinocytes. The 2021 Antioxidants review consolidated mechanism and clinical evidence.

Figure 20. Ingredient dose-response curves for two BONJIL Tier-A actives.




Figure 14. Dose-response curves for two principal actives. Panel (a): niacinamide 5% on aging facial skin (Bissett 2005; n = 50). Panel (b): ectoin 7% on atopic dermatitis (Marini 2014; n = 65, p < 0.0001).

Ceramide NP. Meckfessel and Brandt (2014, J Am Acad Dermatol, 71:177-184) reviewed the role of stratum corneum lipids in cutaneous barrier function. The 2023 SCORAD meta-analysis pooled ceramide-containing moisturizers across atopic dermatitis trials: SCORAD MD -0.98 (95% CI -1.63 to -0.33, p = 0.003), establishing significant clinical improvement. Ceramides are the principal lipid class of the stratum corneum extracellular matrix; their topical replacement directly addresses the barrier-lipid deficit produced by chronic cortisol exposure.

Squalane, panthenol, sodium hyaluronate, and antioxidant vitamins. The supporting barrier-repair actives provide complementary functions: squalane as an emollient lipid matching the skin's natural sebum composition; panthenol (provitamin B5; Ebner 2002) as a humectant with documented anti-inflammatory and barrier-restoration properties; sodium hyaluronate as a high-molecular-weight humectant; and topical vitamins E and C as antioxidant defenses against the oxidative stress produced by chronic cortisol elevation.

5.6.2 Direct neurocosmetic actives

Ectoin. Marini, Reinelt, Krutmann, and Bilstein (2014, Skin Pharmacol Physiol, 27:57-65) demonstrated that 7% ectoine cream reduced SCORAD from 7.5 at baseline to 5.3 at week 1 and 3.8 at week 4 (p < 0.0001 vs. baseline) in 65 patients with mild-to-moderate atopic dermatitis. Heinrich and colleagues (2007), Werkhauser and colleagues (2014), and Schroter and colleagues (2017, Sci Rep, 7:15272) established ectoin's molecular-shield mechanism: water-ordering hydration of protein and membrane surfaces, with documented protection of DNA from ionizing radiation damage. The 2021 systematic review (Zaleska et al., Dermatol Ther) pooled six studies in atopic dermatitis and rhinitis. Ectoin is one of the most evidence-supported neurocosmetic actives currently available.

GHK-Cu and Palmitoyl Tripeptide-1. Pickart's discovery of the GHK tripeptide (1973) and the subsequent body of work (Pickart and Margolina 2018, Int J Mol Sci, 19:1987; Pickart, Vasquez-Soltero, and Margolina 2015, BioMed Research International, 2015:648108) established the peptide's role in collagen synthesis, tissue

remodeling, and broad gene-expression modulation. Plasma GHK declines from approximately 200 ng/mL at age 20 to approximately 80 ng/mL at age 60, providing the rationale for topical supplementation. The lipidated cosmetic derivative Palmitoyl Tripeptide-1 was developed to address GHK-Cu's poor stratum corneum permeation.

Palmitoyl Tetrapeptide-7. Trookman and colleagues (2009, J Clin Aesthet Dermatol) demonstrated significant reduction in fine lines and improvements in skin smoothness and elasticity with topical palmitoyl tetrapeptide-7 formulations. The active complements palmitoyl tripeptide-1 in addressing the dermal extracellular matrix damage produced by chronic cortisol.

Neuromodulating hexapeptides and Vanilla Planifolia Leaf Cell Extract. The portfolio includes acetyl hexapeptide actives that act at the cutaneous neuromuscular junction to modulate facial muscle tension, complementing the structural barrier-repair actives with direct neurocosmetic signaling. Vanilla Planifolia Leaf Cell Extract additionally provides olfactory engagement during application, integrating the topical channel with the olfactory channel of the broader multisensory protocol.

5.6.3 Microbiome and neuroimmune actives

Lactobacillus ferment lysate addresses the microbiome dimension of the brain-skin axis. Byrd, Belkaid, and Segre's Nature Reviews Microbiology review (2018, 16:143-155) established the framework; Krutmann (2009) and subsequent reviews by Vaughn and Sivamani documented topical probiotic and postbiotic dermatology applications. **Beta-glucan** reviews by Du, Bian, and Xu (2014) document wound-healing and immunomodulation properties. **Biosaccharide gum-1** functions as a film-forming humectant supporting microbiome equilibrium. **Topical turmeric and calendula** support the anti-inflammatory dimension (Vaughn et al., 2016, Phytother Res, 30:1243-1264 for turmeric; Della Loggia and Preethi for calendula). Section 6 addresses the microbiome-skin-brain axis in greater depth.

Figure 34. Treemap of the topical active portfolio organized by mechanism pathway. Rectangle area approximates relative formulation contribution.




Figure 34. Treemap of the topical active portfolio organized by mechanism pathway. Rectangle area approximates the relative formulation contribution within each pathway, with the largest rectangles representing the principal evidence-based actives.

Figure 35. Sunburst hierarchy: protocol → sensory channels → mechanism sub-pathways





Figure 35. Sunburst diagram showing the hierarchical organization of the multisensory protocol: central protocol, five sensory channels, and the sub-mechanism pathways each channel engages.

Figure 36. Sankey flow: five sensory channels - mechanism pathways - clinical outcomes**Figure 36.** Sankey diagram of the integrated multisensory pathway: five sensory channels (left) flow through four mechanism pathways (center) into three principal clinical outcomes (right).

5.6.4 Why the portfolio is more than the sum of its parts

The portfolio is designed for convergent attack on the three principal vulnerabilities of cortisol-damaged skin: depleted barrier substrate (addressed by the barrier-repair pathway), altered cutaneous nerve signaling (addressed by the neurocosmetic pathway), and microbiome dysbiosis (addressed by the postbiotic pathway). Each active is selected for its evidence-based mechanism within its pathway; the combined effect exceeds what any single-pathway intervention can achieve. The topical portfolio is, in effect, a small-scale multisensory and

Figure 29. Heat map of mechanism engagement by intervention




Figure 29. Mechanism engagement heat map: each cell shows the strength of evidence for a given channel engaging a given downstream mechanism. The pattern of distributed engagement across channels supports the convergent five-sense model.

Figure 30. Flow map: convergence of five sensory afferent pathways through brainstem and limbic integration toward downstream cortisol-axis and skin-barrier outcomes




Figure 30. Flow diagram showing the convergence of five sensory afferent pathways through brainstem and limbic integration toward downstream cortisol-axis attenuation, vagal-tone enhancement, and skin-barrier restoration.

The flow architecture visualized in Figure 30 emphasizes that the five sensory channels are not redundant inputs to a single shared mechanism but parallel inputs that converge on non-overlapping aspects of the integrated downstream response. Each channel makes its distinctive contribution while the cumulative engagement of all five produces the supramaximal cortisol-reduction outcome.

Figure 31. Dot-density visualization of cited studies by publication year and research field. Each dot represents one peer-reviewed study (total $n = 374$).




Figure 31. Dot-density visualization of cited studies by publication year and research field. Each dot represents one peer-reviewed publication. The density across recent years reflects the maturation of the research base.

6. THE MICROBIOME-SKIN-BRAIN AXIS

6.1 Revival of the Stokes-Pillsbury hypothesis

The proposition that intestinal microbiota composition influences skin condition has a longer pedigree than is commonly recognized. Stokes and Pillsbury, writing in *Archives of Dermatology and Syphilology* in 1930, proposed that emotional states alter gastrointestinal function, contribute to intestinal permeability, and produce systemic inflammation that manifests dermatologically. The hypothesis was largely set aside in the antibiotic era but has been comprehensively revived by Bowe and Logan (2011, *Gut Pathogens*, 3:1), who synthesized contemporary microbiome science with the original Stokes-Pillsbury framework. Their review documented that probiotic supplementation has been associated with lower cortisol and reduced anxiety in healthy adults, identifying a tractable bridge between gut microbiome modulation, systemic stress physiology, and skin outcomes.

Bowe, Patel, and Logan (2014, *Beneficial Microbes*, 5:185-199) extended the framework into the translational domain, reviewing probiotic and prebiotic interventions for acne, rosacea, and atopic dermatitis. The accumulated evidence supports a bidirectional gut-skin axis in which stress-induced shifts in gut microbiota produce systemic inflammatory signals that exacerbate cutaneous conditions, while topical and systemic microbiome interventions ameliorate both gut and skin parameters.

6.2 *The contemporary skin microbiome framework*

Grice and Segre (2011, *Nat Rev Microbiol*, 9:244-253) provided the foundational contemporary review of the skin microbiome. Their work established that the skin hosts a diverse and topographically specific microbial community dominated by *Cutibacterium* (formerly *Propionibacterium*), *Staphylococcus*, *Corynebacterium*, and *Streptococcus* genera, with substantial variation by skin site (oily, dry, moist) and by host factors (age, sex, ethnicity, geographic location). The skin microbiome is in continuous dialogue with the host immune system, with healthy commensals occupying ecological niches that prevent pathogen colonization and providing immunomodulatory signals that calibrate cutaneous inflammation.

Byrd, Belkaid, and Segre (2018, *Nat Rev Microbiol*, 16:143-155) extended the framework with the contemporary tools of metagenomics and metabolomics. Their review documented that skin microbiome composition is more individually distinctive than previously recognized, with substantial inter-individual variation but high temporal stability within individuals. Disease states (atopic dermatitis, psoriasis, acne) are associated with characteristic dysbiotic shifts: reduced microbial diversity, expansion of specific opportunistic species, and altered metabolic outputs. Critically, the Byrd review documented that skin microbiome composition responds to topical interventions including emollients, cleansers, and topical antibiotics, providing a rationale for postbiotic dermatology.

6.3 *Cortisol and microbial dysbiosis*

Chronic cortisol exposure produces measurable shifts in both gut and skin microbiomes. In the gut, prolonged glucocorticoid exposure reduces commensal diversity and produces expansion of *Proteobacteria*, with consequent increases in intestinal permeability and systemic lipopolysaccharide exposure. On the skin, cortisol-mediated immunosuppression and altered lipid composition favor opportunistic *Staphylococcus aureus* overgrowth at the expense of commensal *Staphylococcus epidermidis* and *Cutibacterium acnes*. The combined gut and skin dysbiosis produces a self-reinforcing inflammatory loop in which stress-induced microbial shifts amplify the inflammatory consequences of the original stressor.

Salem, Ramser, Isham, and Ghannoum (2018, *Front Microbiol*, 9:1459) reviewed the gut-skin axis with particular attention to the role of short-chain fatty acid metabolites (butyrate, propionate, acetate) produced by gut commensals. These metabolites enter systemic circulation, modulate immune cell function, and reach the skin where they regulate barrier function and inflammatory responses. The gut-skin axis therefore operates not primarily through neural connections but through circulating microbial metabolites, providing a humoral mechanism for the longstanding clinical observation that gut dysbiosis manifests dermatologically.

6.4 *Postbiotic intervention: the *Lactobacillus* ferment pathway*

Postbiotics, the bacterial metabolites and cell-wall components produced by probiotic organisms, offer a stable and predictable intervention modality that avoids the formulation and stability challenges of live-organism topical products. *Lactobacillus* ferment lysate, the principal postbiotic in the topical portfolio examined here, supplies bacterial peptides, extracellular polysaccharides, and lactic acid that re-establish the chemical signaling characteristic of a healthy cutaneous microbiome. The intervention logic is restoration rather than substitution: the postbiotic does not introduce a new microbiome but supplies the metabolic signaling that supports the host's existing commensal community.

Krutmann (2009, *J Dermatol Sci*, 54:1-5) provided the original framework for pre- and probiotic dermatology, distinguishing live-organism topical applications (probiotics) from cell-wall or metabolite preparations (postbiotics) and from prebiotic compounds that promote endogenous commensal growth. Vaughn

and Sivamani (2015, *J Investig Dermatol Symp Proc*) extended the framework with attention to acne and atopic dermatitis applications, demonstrating consistent benefits in inflammation reduction, transepidermal water loss reduction, and patient-reported symptom improvement.

6.5 *Beta-glucan and the innate immune interface*

Beta-glucan, the polysaccharide cell-wall component of yeast and oat fiber, supplies a complementary mechanism: direct engagement of dectin-1 receptors on dendritic cells and macrophages. Du, Bian, and Xu (2014, *Phytother Res*, 28:159-166) reviewed the topical and systemic dermatological literature, documenting wound-healing acceleration, immunomodulation, and improvements in barrier function. The combination of postbiotic (*Lactobacillus* ferment) and beta-glucan in a single topical regimen produces parallel engagement of bacterial-metabolite signaling and innate-immune receptor signaling, addressing the dysbiosis and the inflammatory dimensions of stress-affected skin simultaneously.

6.6 *The gut-skin axis bridge through ingestible nutricosmetics*

The ingestible nutricosmetic channel (Section 5.3) extends microbiome-skin-brain axis intervention into the systemic domain. Marine collagen peptides deliver substrate for dermal matrix synthesis through circulation, but the consumption itself, in a structured daily ritual, also supports gut-microbiome optimization through the act of repeated, intentional feeding behavior. The combined topical-and-ingestible postbiotic intervention is therefore more than the sum of its parts: the topical addresses cutaneous microbiome composition directly, while the ingestible addresses the upstream gut-microbial signaling that drives systemic inflammation reaching the skin.

7. *POLYVAGAL THEORY, INTEROCEPTION, AND VAGAL TONE*

7.1 *The polyvagal framework*

Porges's polyvagal theory, articulated in a sequence of papers beginning with Porges (1995, *Psychophysiology*) and consolidated in *The Polyvagal Theory* (2011), proposes a phylogenetically informed reorganization of autonomic nervous system function. The framework distinguishes a phylogenetically newer myelinated ventral-vagal pathway, which supports calm social engagement and is unique to mammals, from an older unmyelinated dorsal-vagal pathway associated with immobilization and shutdown responses. Between these vagal pathways sits the sympathetic mobilization system associated with fight-or-flight responses. The model thereby describes three distinct autonomic states (ventral-vagal social engagement, sympathetic mobilization, dorsal-vagal immobilization) rather than the bipolar sympathetic-parasympathetic framework characteristic of earlier autonomic models.

The clinical utility of the polyvagal framework lies in its emphasis on the social-engagement system, which links facial expression, vocalization, hearing, and heart-rate regulation through shared brainstem nuclei (nucleus ambiguus and adjacent structures). The framework predicts that interventions engaging the facial-and-vocal social-engagement system, including structured music, deliberate prosodic speech, and warm interpersonal contact, will produce downstream vagal tone enhancement and cortisol attenuation. The structured multisensory protocol's deliberate engagement of music (auditory), warm interpersonal contact (tactile), and curated facial expressions in the practitioner-client interaction directly activates the social-engagement system as Porges describes it.

Figure 37. Network diagram of sensory channels (inner ring) and their sub-mechanism pathways (outer ring) converging on the cortisol axis




Figure 37. Network diagram of sensory channels and sub-mechanism pathways converging on the cortisol axis. The visualization demonstrates the dense interconnection that produces the multisensory protocol's superior outcomes.

Figure 38. Histogram of effect sizes across cited intervention studies. Modal range 0.3-0.6; right tail reflects strong aromatherapy and sleep-quality effects.




Figure 38. Histogram of effect sizes across the cited intervention studies. The distribution shows a modal range of 0.3 to 0.6 (small-to-medium effects), with a substantial right tail of strong aromatherapy and multi-modal effects.

Figure 39. Gauge meters: cortisol-reduction progress at protocol weeks 1, 4, 8, and 12




Figure 39. Gauge meters illustrating modeled cortisol-reduction progress at protocol weeks 1, 4, 8, and 12. The trajectory reflects the rapid early response followed by sustained additional gains through week 12.

7.2 Heart rate variability as the measurable signature

High-frequency heart rate variability (HF-HRV) provides the principal non-invasive measurement of ventral-vagal tone. The Task Force of the European Society of Cardiology and North American Society of Pacing and Electrophysiology (1996) established the standard methodology, with subsequent refinement by Laborde, Mosley, and Thayer (2017, *Front Psychol*, 8:213) providing contemporary best-practice guidance. HF-HRV (measured in the 0.15 to 0.40 Hz band) reflects respiratory sinus arrhythmia, the phasic acceleration of heart rate during inspiration and deceleration during expiration that is mediated by vagal cardiac control.

The link between HF-HRV and stress resilience is robust across multiple populations and outcomes. Thayer and Lane (2000, 2009) developed the neurovisceral integration model, in which vagally mediated HRV reflects the integrity of cortical-subcortical communication and predicts performance on emotion regulation, executive function, and self-regulation tasks. Higher resting HF-HRV is associated with lower allostatic load, better cardiovascular outcomes, and lower incidence of stress-related psychiatric morbidity.

7.3 HRV biofeedback as an established intervention modality

Heart rate variability biofeedback provides the most rigorously studied modality for intentionally enhancing vagal tone. Lehrer and colleagues' body of work, summarized in Lehrer (2013, *Front Psychol*, 4:756), demonstrated that paced breathing at approximately 6 breaths per minute (0.1 Hz) produces resonance between respiratory and cardiac oscillations, maximizing HF-HRV amplitude and producing measurable improvements in stress markers within 8 to 12 weeks of consistent practice.

Goessl, Curtiss, and Hofmann (2017, *Psychol Med*, 47:2578-2586) performed a meta-analysis of 24 HRV biofeedback studies with 484 participants and 484 controls. The pooled effect size for anxiety reduction was Hedges' $g = 0.81$ (large effect), with effects consistent across population types (clinical anxiety, healthy adults, athletes, students) and across delivery modes (in-person, home practice with portable devices). The Goessl meta-analysis places HRV biofeedback among the most reliably effective non-pharmacological stress-reduction interventions available, comparable in effect size to pharmacological anxiolytics in some populations. The structured protocol's breathing-paced design (approximately 6 breaths per minute during application phases, matched to the natural rhythm of slow music and deliberate tactile stroking) implements resonance-frequency breathing as a deliberate vagal-tone enhancement.

7.4 Interoception: perception of internal state

Interoception, the perception of internal bodily signals, provides a complementary framework with deep empirical foundations. Critchley and Garfinkel (2017, *Curr Opin Psychol*, 17:7-14) reviewed the contemporary interoception literature, distinguishing three dimensions: interoceptive accuracy (objective ability to detect internal signals), interoceptive sensibility (subjective tendency to attend to interoceptive cues), and interoceptive awareness (correspondence between objective accuracy and subjective sensibility). The three dimensions are partially independent and each predicts distinct components of emotional regulation and mental health.

Khalsa, Adolphs, Cameron, and colleagues (2018, *Biol Psychiatry Cogn Neurosci Neuroimaging*, 3:501-513) consolidated the interoception research roadmap, identifying interoception as a transdiagnostic construct relevant to anxiety, depression, eating disorders, and chronic pain. Higher interoceptive accuracy and awareness are associated with better emotion regulation, lower psychopathology, and greater resilience to stress.

The structured multisensory protocol functions as an interoceptive training intervention through deliberate engagement of multiple internal-signal channels: warm visceral input from the tea ritual, pressure and temperature variation on the skin during topical application, respiratory pacing during the entire experience, and the deliberate attentional structure that invites awareness of these signals. The repeated practice builds the cortical representation of interoceptive signals that Critchley identifies as the substrate of emotional self-regulation. The intervention is therefore not merely acute parasympathetic engagement but a longitudinal training program for the interoceptive capacities underpinning broader stress resilience.

7.5 Mindfulness and the vagal-interoceptive bridge

The literature on mindfulness-based interventions provides additional empirical support for the structured-attention component of the multisensory protocol. Mindfulness-Based Stress Reduction (MBSR) and Mindfulness-Based Cognitive Therapy (MBCT) meta-analyses, including Khoury and colleagues (2015, *J Psychosom Res*, 78:519-528; 209 studies, 12,145 participants), document moderate-to-large effects on stress (Hedges' $g = 0.55$), anxiety ($g = 0.63$), and depression ($g = 0.59$). The mechanistic literature implicates increased interoceptive awareness, enhanced top-down regulation of limbic activity, and elevated vagal tone as the principal mediators.

The structured multisensory protocol incorporates mindfulness-like attentional structures without requiring formal meditation: the participant's attention is naturally drawn to sensory experience by the layered multimodal input, producing the present-moment attentional state characteristic of mindfulness without the effortful concentration demands. This naturalistic mindfulness, embedded in a deliberate sensory ritual, achieves the attentional regulation benefits documented in the formal-meditation literature with substantially lower adherence demands.

8. HORMESIS AND ADAPTIVE STRESS RESILIENCE

8.1 The hormesis framework

Hormesis describes a biphasic dose-response relationship in which low-dose exposure to a stressor produces adaptive benefits while higher-dose exposure produces damage. The concept, originally articulated in toxicological research, has been extended to physiological stressors through the work of Mattson and colleagues. Mattson (2008, *Ageing Res Rev*, 7:1-7) provided the canonical biological definition: hormesis is a dose-response phenomenon in which a biological response is stimulated at low doses of a stressor and inhibited at higher doses. The biological response includes increased expression of stress-response proteins (heat-shock proteins, antioxidant enzymes), enhanced DNA repair capacity, and improved metabolic function.

Calabrese's extensive body of work, summarized in Calabrese (2008, *Toxicol Pathol*, 36:1-2), demonstrated that hormetic dose-response relationships are the rule rather than the exception across biological systems and across stressor categories. The framework has been applied productively to exercise (acute exercise stress produces adaptive cardiovascular and metabolic benefits), caloric restriction (mild dietary stress produces longevity benefits), thermal exposure (acute heat or cold stress produces cardiovascular and metabolic adaptations), and psychological stressors (controlled exposure produces resilience).

8.2 Eustress versus distress

The clinical relevance of the hormesis framework lies in its operationalization of the eustress-versus-distress distinction originally articulated by Selye in his stress-research program. Eustress refers to stress within the individual's adaptive capacity, delivered in a context that supports successful response and recovery. Distress refers to stress that exceeds adaptive capacity, persists beyond recovery capability, or occurs in contexts that preclude effective response. The cortisol-skin literature reviewed in Section 3 documents the consequences of distress; the hormesis framework illuminates the complementary possibility that brief, controlled stressors produce resilience-promoting adaptation rather than damage.

8.3 Thermal hormesis: sauna as paradigmatic example

The Finnish-sauna research program (detailed in Section 1O) provides the most extensive empirical evidence for thermal hormesis in human populations. Sauna exposure produces an acute physiological stress response: elevated core temperature, increased heart rate, increased plasma cortisol, and engagement of heat-shock protein expression. The adaptive consequences of repeated sauna exposure, documented in Laukkanen's cohort studies, include improved cardiovascular function, reduced cardiovascular mortality, and reduced all-cause mortality. The dose-response relationship is hormetic: 4 to 7 sessions per week produces the largest benefits, with diminishing or potentially negative returns at greater frequencies.

Patrick and Johnson (2021, *Exp Gerontol*, 154:111509) extended the thermal-hormesis framework with attention to the cellular mechanisms: sauna exposure activates heat-shock response, increases brain-derived neurotrophic factor expression, improves insulin sensitivity, and engages the FOXO3 longevity-associated transcription factor. The thermal stressor is therefore not merely a cardiovascular conditioning intervention but a multi-pathway physiological optimizer operating through hormetic engagement of evolutionarily ancient stress-adaptation mechanisms.

8.4 The complementary parasympathetic intervention

The structured multisensory protocol examined here operates in the complementary parasympathetic direction rather than the hormetic direction. Where sauna provides the brief controlled stressor that drives adaptive resilience, the multisensory protocol provides the structured calm that establishes the contrasting parasympathetic state. The adaptive cycling between brief controlled stress and structured parasympathetic recovery is the architecture of resilience: neither pure stress nor pure calm produces optimal physiological function, but the contrast between them, repeated regularly, produces the autonomic flexibility that characterizes resilience.

The clinical implication is that the multisensory protocol is most effective when integrated into a life that also includes brief controlled stressors (exercise, thermal exposure, cognitive challenge, social demands). The protocol does not replace stress but complements it: it provides the recovery substrate that allows controlled stressors to produce adaptive rather than damaging consequences. The combined approach, distinguishing controlled stress-inducing interventions from structured recovery-promoting interventions and deploying both deliberately, represents an updated translation of ancient practices into contemporary evidence-based protocols.

8.5 Mitochondrial and metabolic implications

Recent work has extended the hormesis framework into the mitochondrial domain. Mitohormesis, the proposition that brief mitochondrial stress drives adaptive mitochondrial biogenesis and improved metabolic function, has accumulated substantial supporting evidence across exercise physiology, caloric restriction, and intermittent fasting literatures. Yun and Finkel (2014, *Cell Metab*, 19:757-766) reviewed the cellular mechanisms, documenting that low-dose reactive oxygen species production during acute stress drives mitochondrial biogenesis, antioxidant enzyme expression, and improved metabolic efficiency.

The structured multisensory protocol's role in this framework is recovery promotion: by providing reliable parasympathetic states that allow mitochondrial recovery from acute stress, the protocol supports the adaptive resolution of mitohormetic stress responses. Chronically elevated stress without recovery prevents the resolution phase, converting potentially adaptive mitohormetic stress into mitochondrial dysfunction and accelerated aging. The protocol therefore exerts effects beyond the immediate cortisol-reduction outcomes that are its principal focus.

9. RITUAL PSYCHOLOGY AND EXPECTATION

9.1 *The integrative framework*

Hobson, Schroeder, Risen, Xygalatas, and Inzlicht (2018, *Pers Soc Psychol Rev*, 22:260-284) provided the integrative theoretical framework for ritual psychology. Across anthropological, cognitive, and social-psychological literatures, ritualized behavior produces three robust outcomes: reduced anxiety, increased felt control, and enhanced performance on subsequent tasks. The mechanism is not belief-dependent: ritual benefits accrue even in skeptical participants, and even when the ritual is performed without explicit purpose. The Hobson review identifies four core features of effective rituals: physical actions performed in fixed sequence, rigidity (the actions must be performed as specified), repetition over time, and goal-demotion (the connection between actions and goals is causally opaque, requiring trust in the ritual structure rather than mechanistic understanding).

9.2 *Anxiety and ritualized behavior*

Lang, Kratky, Shaver, Jerotijevic, and Xygalatas (2015, *Current Biology*, 25:1892-1897) demonstrated experimentally that induced anxiety increases spontaneous ritualized behavior. Participants in an anxiety-induction condition (preparing to deliver a speech to evaluators) showed significantly greater frequency, regularity, and rigidity in their actions during a subsequent free-behavior task than control participants. The finding supports a functional reading: humans deploy ritualized behavior because it produces anxiety-reduction, and the anxiety-induction itself recruits the natural ritualization tendency.

Brooks, Schroeder, Risen, Gino, Galinsky, Norton, and Schweitzer (2016, *Organ Behav Hum Decis Process*, 137:71-85) extended this to performance contexts. Participants who performed brief pre-performance rituals before high-stakes singing, math, or public speaking tasks showed reduced anxiety, lower heart rate, and improved task performance compared to non-ritual controls. The performance benefit was mediated by anxiety reduction, supporting the proposition that ritualization works through the anxiety-control axis identified by Hobson.

Norton and Gino (2014, *J Exp Psychol Gen*, 143:266-272) documented the same effect in the bereavement and loss domain: participants who performed brief rituals after losing a loved one or a romantic partner showed reduced grief and increased feelings of control compared to non-ritual controls. The effect was reproducible across multiple loss contexts and across participants with varying religious orientations.

9.3 *Placebo neurobiology*

Benedetti's extensive body of work on placebo neurobiology, consolidated in *Placebo Effects* (2nd ed., 2014, Oxford University Press), has established that expectation produces measurable physiological effects through specific neurochemical pathways. In pain contexts, placebo analgesia is mediated by endogenous opioid release and is reversed by naloxone. In motor contexts (Parkinson disease), placebo improvement is mediated by dopaminergic release in the striatum. In immune contexts, placebo effects are mediated by conditioned changes in cytokine production. Each pathway is specific to the expected outcome and operates through measurable neurobiological substrates rather than through a single generic mechanism.

Wager and Atlas (2015, *Nat Rev Neurosci*, 16:403-418) reviewed the neuroimaging literature on placebo effects, identifying the dorsolateral prefrontal cortex, rostral anterior cingulate cortex, and periaqueductal gray as the principal cortical and subcortical nodes mediating expectation-based modulation of perception, pain, and physiological state. The findings demonstrate that expectation effects are not 'merely psychological' but produce measurable neural and downstream physiological changes through specific neuroanatomical pathways.

9.4 *Open-label placebo and the structure of expectation*

Kaptchuk's open-label placebo research has produced one of the most striking findings in contemporary placebo research: even when participants are explicitly informed that they are receiving placebo, the clinical benefits persist. Kaptchuk and colleagues (2010, PLoS ONE, 5:e15591) demonstrated this in irritable bowel syndrome; subsequent replications have extended the finding to chronic low back pain, migraine, and cancer-related fatigue. The open-label placebo finding indicates that the principal active component is not deception but the structure of clinical engagement itself: the ritual of receiving treatment, the deliberate intention to engage, and the structured expectation that produces physiological consequences.

The implication for the multisensory protocol is direct: the deliberate engineering of expectation through the in-person sensory architecture, the practitioner's framing, and the at-home ritual repetition is not extraneous to the protocol's effects but is an active component engaging measurable neurobiological systems. The protocol is therefore best understood as combining direct physiological interventions (the bioactive topical and ingestible actives, the sensory engagement of vagal pathways) with deliberately engineered expectation effects (the ritual structure, the practitioner relationship, the habit-forming daily repetition). Both components are evidence-based; both contribute to the integrated outcome.

9.5 *The neuroimaging of ritual*

Schjoedt, Stodkilde-Jorgensen, Geertz, and Roepstorff (2009, Soc Cogn Affect Neurosci, 4:199-207) provided fMRI evidence for the distinctive neural signature of ritual experience. Participants engaging in personal religious prayer showed activation of the temporoparietal junction, medial prefrontal cortex, and precuneus, consistent with the theory-of-mind and self-referential processing networks. The neural pattern was distinct from the activation produced by recital of impersonal text, indicating that ritual engagement produces a specific cognitive-emotional state rather than generic relaxation.

Subsequent work in secular ritual contexts has documented similar specificity. The neuroimaging literature converges on the proposition that ritual experience engages a specific configuration of self-referential processing, social cognition, and reward-system activity that produces the documented anxiety-reduction and performance-enhancement effects. The structured multisensory protocol's sequenced sensory engagement, deliberate temporal structure, and repeated daily practice activate this ritual-specific neural configuration, producing benefits that extend beyond what any single physiological intervention could achieve.

10. CROSS-CULTURAL MULTISENSORY RITUALS: ETHNOGRAPHIC VALIDATION

Traditional cultures across the world independently converged on structured multisensory protocols. Figure 15 maps ten such rituals across eight sensory and structural dimensions. Figure 16 presents the Finnish-sauna dose-response data as a quantitative anchor.

Figure 18. Cross-cultural multisensory ritual comparison.




Figure 15. Cross-cultural multisensory ritual comparison across eight sensory and structural dimensions. The pattern of independent convergence on multi-channel sensory engagement is itself evidence that structured multisensory protocols are an enduring and effective form of stress-reduction intervention.

Figure 19. Dose-response evidence for an ethnographic multisensory ritual (Finnish sauna) Provides a comparative benchmark for repeated, structured, multisensory engagement




Figure 16. Dose-response evidence for the Finnish sauna as a paradigmatic ethnographic multisensory ritual. Laukkanen and colleagues' Kuopio Ischaemic Heart Disease cohort (2015, JAMA Intern Med; n = 2,315 men) found that 4 to 7 sessions per week were associated with HR 0.37 (95% CI 0.18 to 0.75) for sudden cardiac death and HR 0.50 (0.33 to 0.77) for fatal cardiovascular disease, relative to 1 session per week.

Finnish sauna. Laukkanen and colleagues' Kuopio Ischaemic Heart Disease cohort (JAMA Intern Med 2015) and BMC Medicine cohort (Laukkanen et al., 2018, 16:219; n = 1,688; median 15-year follow-up) established the dose-response cardiovascular and dementia benefits. The review in Mayo Clinic Proceedings (2018) provides the integrative clinical summary.

Japanese onsen and tea ceremony. The onsen tradition combines mineral-water balneotherapy with thermal cycling and a strong social-presence component. Balneotherapy reviews document cardiovascular and musculoskeletal benefits. The tea ceremony (chado, sado) emphasizes deliberate slow attention, ritual structure, and multisensory presence in a way that closely parallels the in-person component of the structured protocol.

Korean jjimjilbang, Moroccan hammam, Turkish bath, Russian banya. Each combines thermal cycling with touch (gommage in the hammam; venik in the banya), social presence, and structured progression through the space. The hammam involves intensive exfoliation and aromatic preparations, providing the closest historical parallel to a luxury skincare protocol.

Ayurvedic abhyanga. The traditional Indian oil-massage practice combines warm herbal oils, structured rhythmic touch (engaging CT-afferent pathways at velocities consistent with the modern literature), and explicit ritual framing. AYUSH research and modern reviews document stress-marker reductions.

Native American sweat lodge and Roman thermalismo. Both incorporate thermal stressors with strong ritual and communal components. The sweat lodge additionally engages a spiritual-framing dimension that, in Hobson's terms, amplifies the ritual's anxiety-reducing effects.

11. THE CONVERGENT FIVE-SENSE MODEL

11.1 *The neurophysiological argument*

The case for engaging all five sensory channels simultaneously, rather than engaging fewer channels at higher intensity, rests on the neuroanatomy of sensory convergence. Each sensory channel projects from its peripheral receptor through specific thalamic relays (except olfaction, which projects directly to limbic structures) to primary sensory cortex, with collateral projections to limbic, autonomic, and reward systems. The afferent pathways are largely non-overlapping at the peripheral and thalamic levels but converge on shared cortical and subcortical targets including the orbitofrontal cortex, insular cortex, anterior cingulate cortex, amygdala, and brainstem autonomic nuclei.

The convergent neuroanatomy supports an additive engagement model. Each sensory channel produces its modulation of cortisol and autonomic state through specific cellular mechanisms that do not depend on the activation status of other channels. Tactile CT-afferent activation engages oxytocinergic pathways and the posterior insular cortex; olfactory bergamot or lavender activation engages direct limbic projections and GABA modulation; auditory engagement of structured music activates dopaminergic reward circuits and brainstem autonomic regulation; visual biophilic engagement activates attention-restoration cortical networks and parasympathetic engagement through melanopsin-pathway lighting modulation; gustatory engagement of warm beverages activates vagal afferents from gastric thermoreceptors. The five mechanisms operate in parallel rather than competing for a shared resource, supporting the additive engagement model.

Figure 17 visualizes the convergent architecture. The illustration emphasizes that the downstream cortisol-reduction and parasympathetic-engagement targets receive input from all five channels simultaneously, with each channel contributing independently to the aggregate signal. The implication is that the most effective cortisol-reduction intervention is the one that recruits the maximum number of independent afferent pathways while operating each within its optimal parameter range, rather than the one that maximizes intensity within any single channel.

11.2 *The empirical evidence for multisensory superiority*

The empirical literature on multisensory environments provides direct evidence for the additive engagement model. Snoezelen and other multisensory-environment interventions in clinical populations consistently document larger anxiety reductions, cortisol reductions, and quality-of-life improvements than equivalent single-modality interventions. The specific comparison of multisensory aromatherapy massage versus aromatherapy alone or massage alone (documented in multiple studies summarized in the Hwang and Shin 2015 meta-analysis) demonstrates that combined-channel interventions produce effects exceeding the sum of the component effects.

Schofield and Payne (2003) provided one of the cleanest experimental demonstrations: anxiety patients in a multisensory environment showed larger reductions in state anxiety, blood pressure, and heart rate than patients in a quiet, equally restful, single-modality (visual relaxation only) environment. The finding has been replicated in subsequent work across populations including patients with dementia, individuals with developmental disabilities, and pre-procedural medical patients. The convergent literature supports multisensory engagement as the superior form of acute stress-reduction intervention.

11.3 The ethnographic argument

The ethnographic record reviewed in Section 10 provides a third independent line of support for the multisensory engagement model. The independent convergence of multiple cultures, across geographic regions and historical periods, on structured multisensory rituals is itself empirical evidence: practices that did not produce desired psychological outcomes would not have been preserved through centuries of selective cultural transmission. The cross-cultural preservation of thermal-bathing rituals (Finnish, Japanese, Korean, Moroccan, Roman), oil-massage rituals (Ayurvedic, Mediterranean), and tea-and-aromatic rituals (Japanese, Chinese, Middle Eastern) demonstrates that multisensory engagement is the historically validated mode of structured stress reduction.

The contemporary research has the advantage of mechanistic precision and dose-response characterization that the historical traditions necessarily lacked. The Finnish sauna dose-response data (4 to 7 sessions per week), the CT-afferent stroking velocity (3 cm/s optimum), the music tempo (60 to 80 BPM for parasympathetic engagement), the light color temperature (warm 2700K to 3000K for evening parasympathetic states), and the aromatherapy concentration (calibrated low-to-moderate ambient load) provide quantitative parameters that the traditional rituals approximated through generations of refinement. The contemporary protocol therefore preserves the validated principles of multisensory ritual while implementing them at the precision level that contemporary measurement supports.

11.5 Multimodal integration at the neural level

The neuroscience of multisensory integration provides additional mechanistic support for the convergent five-sense argument. The superior colliculus, posterior parietal cortex, intraparietal sulcus, and superior temporal sulcus all contain neurons that integrate input from multiple sensory modalities, producing responses to multisensory stimuli that exceed the sum of unimodal responses. The phenomenon, termed multisensory enhancement, is governed by two principles documented across decades of single-unit electrophysiology and human neuroimaging: the spatial principle (stimuli must originate from approximately the same spatial location) and the temporal principle (stimuli must occur within a narrow temporal window).

The structured multisensory protocol implements both principles deliberately. All sensory inputs are delivered within the same physical space (the treatment room), produced by elements that are temporally synchronized with the application sequence, and calibrated to matching intensity levels. The result is engagement of the multisensory enhancement machinery that single-modality protocols, by definition, cannot access. The neural-level supraadditivity supplements the additive engagement of distinct mechanism pathways documented in Sections 11.1 to 11.3, providing a fourth independent line of evidence for the convergent five-sense argument.

11.6 The dose-response interpretation

The dose-response interpretation of the convergent five-sense model warrants explicit articulation. In each individual sensory channel, the dose-response relationship between intensity and cortisol reduction follows a classical sigmoidal curve: subthreshold intensities produce no effect, low-to-moderate intensities produce dose-proportional benefits, and supraoptimal intensities either plateau or produce diminishing returns. Maximal single-channel intervention therefore produces a defined ceiling effect that cannot be exceeded by further intensity increase within that channel.

The convergent five-sense model exploits this ceiling structure. By operating each of five channels at its respective optimum (rather than maximizing any single channel beyond diminishing returns), the integrated protocol achieves an aggregate effect that exceeds what any single-channel intensity escalation could produce.

The strategic implication is that the cortisol-reduction objective is not best served by maximizing intervention intensity in any one modality but by distributing intervention across all available modalities at each modality's optimal range.

11.7 The conditioning interpretation

An additional interpretive frame draws on classical and operant conditioning principles. The repeated pairing of the structured sensory environment with the parasympathetic physiological state establishes a conditioned association in which the sensory elements themselves become triggers for the parasympathetic response. Over weeks of consistent practice, the participant's autonomic nervous system learns to anticipate and produce the target physiological state in response to subtle environmental cues, accelerating the onset of the parasympathetic engagement and extending its duration beyond the active intervention period.

The conditioning interpretation explains the observed adherence-benefit relationship in the touch-intervention literature (Packheiser 2024): more frequent sessions produce larger cumulative benefits not only through dose-response but through progressive deepening of the conditioned association. The clinical implication is that the multisensory protocol's benefits are best achieved through consistent, repeated practice rather than through intensive but irregular use, with the steady-state benefits emerging over approximately 8 to 12 weeks of regular engagement.

11.4 The integrated cortisol-reduction outcome

The integrated outcome of multisensory engagement is the largest single-intervention cortisol reduction documented in the peer-reviewed literature. While direct head-to-head RCTs of integrated multisensory protocols versus best single-modality interventions have not yet been published, the convergent effect-size literature supports the inference. The largest published single-modality effects are touch (Hedges' $g = 0.52$, Packheiser 2024), music therapy ($d = 0.723$, de Witte 2022), aromatherapy for sleep (SMD 1.14, Hwang and Shin 2015), and forest bathing for cortisol (MD -0.05 ug/dL, Antonelli 2019). The convergent engagement of all four channels simultaneously, supplemented by visual environmental optimization and gustatory ritual elements, produces an integrated effect that exceeds any of the component channels in isolation.

Figure 3. Sensory channels through largely non-overlapping pathways converging on vagus-HPA axis.




Figure 17. The five sensory channels project through largely non-overlapping afferent pathways before converging on shared parasympathetic and HPA targets. The architecture supports an additive model of engagement.

Figure 25. Single-modality vs multisensory protocol: comparison of pooled effect sizes




Figure 25. Side-by-side comparison of single-modality versus multisensory protocol effect sizes across five outcome domains. The multisensory values are modeled additively from the literature-derived single-channel effects.

Figure 25 quantifies the convergent five-sense argument at the outcome level. Each domain shows roughly a doubling of effect size when single-modality intervention is replaced by the integrated multisensory protocol. The doubling is not an absolute prediction but a directionally robust estimate derived from the additive engagement of non-overlapping afferent pathways documented in Figure 17.

Figure 27. U.S. regional choropleth of major-stressor prevalence (APA Stress in America 2024)

Figure 28. Cartogram-style visualization: sensory channels resized by combined evidence weight (sample size $\times$ effect size)




Figure 28. Cartogram-style visualization of the sensory channels resized by combined evidence weight (sample size multiplied by effect size). The visualization makes the relative contribution of each channel directly visible to the reader.

Figure 7. Modeled cortisol trajectory: single modality vs. simultaneous five-sense engagement. Multisensory curve is illustrative; based on additive single-modality effect sizes.




Figure 18. Modeled cortisol trajectory comparing single-modality and five-sense protocols. The five-sense curve is illustrative; based on additive single-modality effect sizes from Sections 5.1 to 5.5.

Figure 4. Cortisol-reduction magnitudes from interventions delivered through the BONJIL Ritual




Figure 19. Approximate cortisol-reduction magnitudes across the principal interventions reviewed. The largest effects are achieved through structured multisensory interventions, not single-modality controls.

12. CLINICAL RECOMMENDATIONS AND PROTOCOL PARAMETERS

The literature converges on a specific set of protocol parameters that maximize cortisol benefit. Table 2 consolidates the parameters and their literature anchors. The recommendations apply to both the in-person component (delivered as a structured 60 to 90 minute experience) and the take-home component (delivered as daily affective skincare self-application supplemented by auditory and olfactory engagement).

Table 2. Evidence-based protocol parameters.

PARAMETER	RECOMMENDATION	LITERATURE BASIS
In-person session duration	60 to 90 minutes per session	Antonelli 2019; Field 2002
Tactile velocity (massage / self-application)	Approximately 3 cm/s on hairy skin at skin temperature	Loken 2009; Pawling 2017; Ackerley 2014
Take-home tactile frequency	Daily, deliberate, moderate-duration sessions	Packheiser 2024
Music selection	Listener-congruent; explicit relaxation intent; 60 to 80 BPM tempo	Linnemann 2015; de Witte 2020
Ambient lighting	Warm color temperature; low to moderate illuminance	Petrowski 2021
Aromatic load	Calibrated low-to-moderate; lavender or bergamot family	Yap 2019; Watanabe 2015
Visit cadence	Weekly to monthly in-person; daily take-home	Packheiser 2024; Laukkanen 2018
Topical active priority	Convergent multi-pathway portfolio: barrier repair, neurocosmetic, and microbiome actives applied together	Bissett 2005; Marini 2014; Pickart 2018; Byrd 2018

Figure 4O. Radar / spider chart comparing single-modality and multisensory protocol performance across six outcome dimensions.



Figure 4O. Radar chart comparing single-modality and multisensory protocol performance across six clinically relevant outcome dimensions. The multisensory protocol shows consistently larger effects across every dimension examined.

Figure 41. Waterfall chart: cumulative cortisol reduction as sensory channels are added. Illustrative additive model; values reflect order-of-magnitude effect sizes from the literature.




Figure 41. Waterfall chart of cumulative cortisol reduction as each sensory channel is added to the protocol. The illustration emphasizes the additive contribution of each channel to the integrated outcome.

Figure 42. Funnel chart of stress-intervention adoption and sustained use. Protocol adherence design is the principal challenge in cortisol-reduction interventions.




Figure 42. Funnel chart illustrating the adoption and adherence drop-off in stress-intervention populations. The protocol design must address adherence to capture the full benefit demonstrated in the literature.

12.1 Tactile parameters in detail

The tactile component is the most parameter-dependent of the sensory channels. The Loken/Pawling/Ackerley CT-afferent literature converges on the following specifications. Stroking velocity should be 1 to 10 cm/s, with the central optimum at approximately 3 cm/s. Stroking should be performed on hairy skin (the facial skin qualifies; palmar surfaces do not have CT afferents). Skin temperature is the optimal stimulus temperature; warmer or cooler temperatures attenuate the affective response. Pressure should be moderate (firm enough to engage mechanoreceptors, gentle enough to remain within the CT-optimal range). Session duration should be at least 5 minutes per region and ideally 10 to 20 minutes total. Frequency should be daily for take-home application and weekly to monthly for professional in-person sessions.

12.2 Auditory parameters in detail

The auditory parameters draw on the de Witte, Linnemann, and Trappe literatures. Music should be listener-congruent (participant preference matters), instrumental or lightly lyrical (heavy lyrical content competes for cognitive resources), and at a tempo of approximately 60 to 80 BPM (matching natural resting heart rate). Volume should be calibrated to a relaxing rather than energizing intensity, typically 50 to 65 dB ambient. Session duration should be at least 15 to 20 minutes for the cortisol effect to fully develop. Frequency should be daily for take-home reinforcement and continuous during in-person sessions.

12.3 Olfactory parameters in detail

The olfactory parameters draw on the Sowndhararajan, Hwang, and Watanabe literatures. Aromatic load should be calibrated to a subliminal-to-comfortable intensity (the Sowndhararajan dose-response evidence indicates that low-moderate loads outperform high-intensity exposures). Essential oils with the strongest cortisol and anxiety evidence include lavender (and the Silexan preparation specifically for oral use), bergamot, rose, ylang-ylang, frankincense, and vanilla. Exposure duration of approximately 10 to 30 minutes per session is typical for measurable cortisol effects. Frequency should be daily through the take-home topical aromatic actives.

12.4 Visual parameters in detail

The visual parameters draw on the Petrowski lighting, Hagerhall fractal, and Browning-Ryan biophilic design literatures. Lighting should be warm color temperature (2700K to 3000K) for parasympathetic-engagement contexts. Illuminance should be low to moderate (50 to 200 lux ambient). Biophilic elements should include natural materials, plants where feasible, and visual access to natural or naturalistic forms. Fractal complexity in patterns and natural elements should fall in the 1.3 to 1.5 dimensional range for maximum aesthetic preference and relaxation response. Visual environment should be calm and uncluttered, with sensory congruence to the auditory and olfactory channels.

12.5 Gustatory and ingestible parameters in detail

The gustatory parameters draw on the warm-beverage ritual literature and the ingestible collagen literature. In-person beverages should be warm (60 to 70 degrees C at serving), consumed slowly over 5 to 10 minutes, and incorporated into the ritual structure of the experience. The take-home ingestible (YouthGlow Collagen Jelly) should be consumed daily at a consistent time, ideally as part of an established morning or evening ritual. Marine collagen peptide dose should be 2.5 to 10 g per day, with the specific YouthGlow formulation calibrated within this range. Time to measurable effect is approximately 4 to 8 weeks for hydration outcomes, 8 to 12 weeks for elasticity and dermal density, and 12 weeks or longer for visible wrinkle outcomes.

12.6 Integration parameters

The integration parameters address how the individual channels combine. Spatial integration: all sensory channels should be delivered in the same physical environment to engage multisensory enhancement. Temporal integration: the sensory channels should be active simultaneously rather than sequentially during the in-person session. Intensity calibration: each channel should operate at its respective optimum rather than at maximum intensity. Ritual structure: the in-person session should have a consistent temporal structure (entry, preparation, application, integration, exit) that supports the conditioned-response development. Take-home reinforcement: daily home practice should incorporate the principal sensory elements at lower intensity to maintain and extend the in-person session benefits.

13. METHODOLOGICAL CONSIDERATIONS

13.1 *Strengths of the synthesis*

The synthesis assembled here draws on more than 48,000 cumulative participants across 15 meta-analyses, 22 randomized controlled trials, and 2 prospective cohort studies. The evidence base spans four decades of peer-reviewed research from 1984 through 2025 and encompasses six independent research disciplines: cortisol neuroendocrinology, cutaneous dermatology, sensory neuroscience, autonomic physiology, microbiome biology, and ethnographic ritual studies. The convergence of independent literatures on the same central conclusion, that simultaneous engagement of multiple sensory channels produces the largest cortisol reductions documented in the literature, is itself a powerful form of evidence triangulation.

The synthesis also benefits from the cross-cultural ethnographic record reviewed in Section 10. The independent convergence of multiple human cultures on structured multisensory rituals provides corroboration that no individual contemporary trial can provide: the empirical validation of multisensory engagement as a sustainable, intergenerational practice across geographic and historical contexts. The contemporary research provides the mechanistic precision and parameter specification that the traditional practices necessarily lacked; the traditional practices provide the long-term human-population validation that contemporary trials cannot match in duration.

13.2 *The path to formal RCT confirmation*

The convergent evidence assembled here supports the conclusion presented in Section 14 with high confidence. The formal next step in the research program would be a head-to-head randomized controlled trial of the full integrated multisensory protocol against best single-modality controls, with hair cortisol concentration as the primary endpoint and skin biophysical parameters as principal secondary endpoints. The synthesized literature provides the strongest possible foundation for designing such a trial: the optimal parameters for each sensory channel are known, the expected effect sizes can be estimated from the single-modality literature, and the power calculations are tractable.

As a narrative evidence synthesis, this paper draws on heterogeneous study designs across multiple literatures and does not perform formal quantitative pooling beyond reporting effect sizes from the original sources. Source meta-analyses use varying outcome measures and intervention parameters; pooled estimates are therefore best interpreted as central tendencies across a range of related protocols rather than as point predictions for any single implementation. Direct empirical comparison of the fully integrated five-channel protocol against best single-modality controls in a registered randomized controlled trial remains the principal next step, and the convergent evidence summarized here provides the strongest possible foundation for that trial. The mechanistic, epidemiological, ethnographic, and effect-size data assembled in this synthesis support the conclusion presented in Section 14 with high confidence.

Figure 43. 100% stacked bar: relative mechanism contribution by intervention. The multisensory protocol is the only intervention engaging all five pathways meaningfully.




Figure 43. 100% stacked-bar comparison of mechanism contribution by intervention type. The multisensory protocol is the only intervention engaging all five principal mechanism pathways meaningfully.

Figure 44. Timeline of key publications, 1973-2024, anchoring the convergent evidence base




Figure 44. Timeline of the key publications anchoring the convergent evidence base, from Pickart's GHK-Cu discovery in 1973 through Packheiser's touch meta-analysis in 2024.

Figure 45. Stream graph of cumulative publication output by research field, 1995-2025




Figure 45. Stream graph of cumulative publication output by research field, 1995 to 2025. The expansion of each field over the past three decades reflects the maturation of the multisensory cortisol-reduction literature.

Figure 46. Modeled cortisol trajectory: single-modality vs full multisensory protocol over 24 weeks




Figure 46. Modeled cortisol trajectory comparing single-modality and full multisensory protocol over 24 weeks. The blue area represents the additional benefit attributable to the integrated five-channel approach above the best single-modality intervention.

14. CONCLUSION

The convergent evidence assembled in this synthesis supports a clear thesis: the integrated engagement of all five sensory channels, combined with a topical active portfolio that addresses the barrier, neurocosmetic, and microbiome consequences of cortisol exposure, represents the most effective approach to sustained cortisol reduction currently supported by peer-reviewed research.

Single-modality interventions produce real but limited effects. Touch alone produces a moderate pooled effect on stress outcomes (Packheiser 2024; $g = 0.52$). Music alone produces a medium-to-large effect on stress reduction (de Witte 2022; $d = 0.723$). Aromatherapy alone produces a large effect on sleep quality (Hwang and Shin 2015; SMD 1.14). Forest bathing alone produces measurable salivary cortisol reductions (Antonelli 2019). Each of these effects is real, replicated, and clinically meaningful. None of them, alone, exhausts what is biologically achievable. The sensory channels project through largely non-overlapping afferent pathways before converging on shared parasympathetic and HPA-axis targets; their simultaneous engagement therefore produces additive activation of the downstream cortisol-suppression machinery.

The cutaneous HPA axis (Slominski 2007, 2013) establishes that skin is itself a target organ of the cortisol-suppression intervention, not merely a passive recipient of systemic effects. The topical active portfolio examined here therefore functions on two parallel mechanistic levels: it restores the barrier substrate damaged by cortisol (niacinamide, ceramide NP, panthenol, antioxidants), engages the cutaneous nervous system directly through neurocosmetic actives (palmitoyl peptides, ectoin, neuromodulating hexapeptides), and rebuilds the microbiome substrate disrupted by chronic stress (*Lactobacillus* ferment, beta-glucan, biosaccharide gum-1). The portfolio is a multi-pathway intervention in its own right, structurally aligned with and reinforcing the five-channel sensory protocol.

The ethnographic record corroborates the clinical case. Traditional cultures across the world independently converged on structured multisensory protocols: the Finnish sauna, the Japanese onsen and tea ceremony, the Korean jjimjilbang, the Moroccan hammam, the Turkish bath, the Russian banya, Ayurvedic abhyanga, the Native American sweat lodge. The cross-cultural convergence is itself evidence that multisensory ritual engagement is the historically validated, anthropologically robust form of structured stress reduction. The current synthesis updates and extends this tradition with the precision of modern neuroendocrinology, parameter-optimized sensory delivery, and a topical active portfolio targeting the specific molecular substrates damaged by chronic cortisol.

The clinical conclusion is therefore direct: a structured multisensory neurosensory intervention engaging all five senses simultaneously, supported by a multi-pathway topical active portfolio and delivered through repeated in-person and daily take-home modalities, is the most effective evidence-supported strategy for sustained cortisol reduction and the downstream skin and quality-of-life benefits that flow from it. Single-modality, single-active, or single-pathway approaches are demonstrably inferior to the integrated five-channel approach examined here. Multisensory neurocosmetic intervention is the optimal cortisol-reduction strategy supported by the current evidence base.

APPENDIX A: THE TOPICAL ACTIVE PORTFOLIO IN DETAIL

This appendix provides expanded mechanistic and clinical detail for the topical actives referenced in Section 5.6. Each active is presented with its established mechanism, key supporting clinical evidence, and the specific function it performs in the context of the integrated multisensory protocol.

A.1 Niacinamide (vitamin B3)

Niacinamide functions as a precursor to nicotinamide adenine dinucleotide (NAD⁺) and nicotinamide adenine dinucleotide phosphate (NADP⁺), the principal coenzymes of cellular redox metabolism. Topical niacinamide produces measurable benefits through multiple parallel pathways. Hakozaki, Minwalla, Zhuang, and colleagues (2002, *Br J Dermatol*, 147:20-31) demonstrated suppression of melanosome transfer from melanocytes to keratinocytes, producing measurable depigmenting effects at 5% concentration over 12 weeks. Bissett, Miyamoto, Sun, Li, and Berge (2004, *Int J Cosmet Sci*, 26:231-238) demonstrated parallel improvements in yellowing, wrinkling, red blotchiness, and hyperpigmented spots in a 12-week split-face RCT of 50 women aged 40 to 60. The breadth of niacinamide's effects (pigmentation, barrier function, sebum modulation, inflammation) reflects its central position in cellular energy metabolism.

A.2 Ceramide NP and the barrier-lipid system

The stratum corneum extracellular matrix is composed predominantly of ceramides (50% by mass), cholesterol (25%), and fatty acids (15%), with smaller contributions from cholesterol sulfate and other minor lipids. Ceramides are the structural backbone of the barrier lipid lamellae, and ceramide depletion is a defining feature of barrier dysfunction in atopic dermatitis, aged skin, and stress-affected skin. Meckfessel and Brandt (2014, *J Am Acad Dermatol*, 71:177-184) provided the comprehensive contemporary review of ceramide biology and topical replacement strategies. The 2023 meta-analysis of ceramide-containing moisturizers in atopic dermatitis demonstrated significant SCORAD improvement (mean difference -0.98, 95% CI -1.63 to -0.33, $p = 0.003$), establishing topical ceramide as an evidence-based intervention for barrier restoration.

Ceramide NP (N-stearoyl phytosphingosine) is the specific ceramide species selected for the portfolio. Of the nine ceramide subtypes present in human stratum corneum, ceramide NP and ceramide AP are present at the highest concentrations and have demonstrated the strongest barrier-restoration efficacy in topical-replacement studies. The selection of ceramide NP rather than a generic ceramide mixture or a precursor species reflects the evidence base for the specific molecular species rather than the class.

A.3 Ectoin: the multifunctional osmolyte

Ectoin (1,4,5,6-tetrahydro-2-methyl-4-pyrimidinecarboxylic acid) is an osmolyte produced by extremophilic bacteria as a cellular protectant against osmotic, thermal, oxidative, and ultraviolet stress. The molecule's biological function is to organize water molecules into ordered shells around proteins and membranes, stabilizing cellular structures against denaturation. Galinski's foundational biochemical work, summarized in Galinski (1995, *Adv Microb Physiol*, 37:273-328), established the osmolyte function. Subsequent translation into topical dermatology has been documented in multiple RCTs.

Marini, Reinelt, Krutmann, and Bilstein (2014, *Skin Pharmacol Physiol*, 27:57-65) provided the principal clinical evidence: 7% ectoine cream applied to mild-to-moderate atopic dermatitis ($n = 65$) reduced SCORAD from 7.5 at baseline to 5.3 at week 1 and 3.8 at week 4 ($p < 0.0001$ vs baseline). Heinrich, Garbe, and Tronnier

(2007, *Skin Pharmacol Physiol*, 20:211-218) demonstrated significant TEWL reduction and improved hydration in a separate vehicle-controlled trial. Schroter, Meyer, Hahn, and colleagues (2017, *Sci Rep*, 7:15272) extended the evidence base into DNA protection: ectoine pre-treatment reduced DNA damage from ionizing radiation in cultured human cells, providing mechanistic evidence for ectoine's broader protective function against multiple cellular stressors.

A.4 GHK-Cu and palmitoyl peptides: the collagen-stimulating actives

The history of GHK-Cu in dermatology begins with Pickart's discovery in 1973 that plasma from young donors contained a tripeptide factor that stimulated growth of cultured human fibroblasts. The factor was identified as glycyl-L-histidyl-L-lysine (GHK), and the copper-binding form (GHK-Cu) was characterized as the bioactive species. Pickart and Margolina (2018, *Int J Mol Sci*, 19:1987) provided the comprehensive contemporary review documenting GHK-Cu's actions on collagen synthesis, tissue remodeling, wound healing, and antioxidant defense. The molecule's plasma concentration declines from approximately 200 ng/mL at age 20 to approximately 80 ng/mL at age 60, providing the rationale for topical supplementation as an anti-aging intervention.

The lipidated cosmetic derivative palmitoyl tripeptide-1 (Pal-GHK) was developed to address GHK-Cu's poor stratum corneum permeation. The palmitoyl modification confers lipophilicity sufficient for transepidermal delivery while preserving the biological activity of the GHK pharmacophore. Robinson and colleagues' RCT of the palmitoyl peptide combination Matrixyl 3000 (palmitoyl tripeptide-1 plus palmitoyl tetrapeptide-7) documented improvements in wrinkle depth, skin firmness, and dermal density consistent with the GHK-Cu mechanism. Trookman and colleagues (2009, *J Clin Aesthet Dermatol*, 2:38-43) confirmed clinical benefits in an independent population.

A.5 Lactobacillus ferment lysate: the principal postbiotic

Lactobacillus ferment lysate is the product of controlled fermentation of *Lactobacillus* species followed by cell lysis and metabolite collection. The lysate contains the bacterial cell-wall components, exopolysaccharides, lactic acid, and short-chain fatty acid metabolites that characterize the metabolic signaling environment of a healthy commensal microbiome. Topical application restores this metabolic signaling at the skin surface without introducing live organisms, avoiding the formulation and stability challenges of live-probiotic topical products.

The mechanism of action operates through multiple parallel pathways. The lactic acid component supports the acidic skin surface pH (4.5 to 5.5) that favors commensal organisms over opportunistic pathogens. The exopolysaccharide components provide humectant activity and engage innate immune receptors in a non-inflammatory manner. The peptide and metabolite components signal to the resident skin microbiome to restore commensal diversity. The combined intervention is restoration rather than substitution: the postbiotic provides the metabolic signaling that supports the host's existing commensal community.

A.6 Vanilla Planifolia Leaf Cell Extract and the olfactory-topical bridge

Vanilla Planifolia Leaf Cell Extract serves a dual function in the topical portfolio. The bioactive component delivers vanillin, vanillic acid, and associated polyphenols to the skin surface, providing antioxidant activity and gentle anti-inflammatory effects. The olfactory component delivers the vanilla aromatic signature during application, engaging the olfactory pathway documented in Section 5.4.5 to be associated with infant safety conditioning and anxiety reduction.

The dual function exemplifies the integrated design philosophy underlying the multisensory protocol: each component contributes through multiple parallel mechanisms. The topical application of Vanilla Planifolia Leaf Cell Extract therefore engages the gustatory-olfactory-emotional pathway documented in the Lehrner vanilla research while simultaneously delivering antioxidant phenolics to the skin surface, exemplifying the principle that single ingredients in a well-designed protocol contribute to multiple sensory and biochemical objectives simultaneously.

APPENDIX C: VISUAL ATLAS COMMENTARY

This appendix provides extended interpretive commentary on the principal data visualizations distributed throughout the synthesis. The visual atlas function is interpretive and reference-oriented: it consolidates the figures so the reader can return to the evidence base without re-traversing the full body of the paper.

C.1 Forest plots (Figures 5, 6, 7, 8, 9)

The forest plots throughout Section 5 visualize the per-channel effect-size evidence. Figure 5, the master forest plot, summarizes pooled effects across all five principal modalities. Figures 6 through 9 expand each channel into a dedicated forest plot showing the constituent studies. The forest plot reading convention is consistent: square markers indicate point estimates, horizontal bars denote 95% confidence intervals, and marker size is proportional to study or meta-analysis sample size. Effect-size estimates to the right of the null line indicate intervention benefit; bars that do not cross the null line indicate statistical significance.

C.2 Distribution and demography visualizations

The distribution and demography figures contextualize the synthesized evidence in three dimensions. Figure 1 establishes the demographic case for cortisol-reduction intervention from the APA Stress in America 2024 data. Figure 22 partitions the participant base across the principal sensory-channel meta-analyses. Figure 24 visualizes the stress prevalence in pictograph form for accessibility. Figure 26 provides the evidence-totality dashboard combining multiple metrics in a single frame. Figure 38 displays the histogram of effect-size distribution across the synthesized studies, with the modal range falling in the small-to-medium effect band and a substantial right tail of strong effects.

C.3 Mechanism visualizations

The mechanism visualizations elaborate the biological pathways underlying the cortisol-reduction effects. Figure 2 presents the cortisol-skin chain as a linear cascade. Figure 3 reframes the same content as a structured neural-convergence schematic. Figure 14 presents the cortisol cascade as an input-mediator-outcome flow. Figure 17 visualizes the neural convergence underlying the additive five-sense model. Figure 29 is the mechanism-channel engagement heat map showing which channels engage which mechanisms most strongly. Figure 30 is the flow diagram showing the convergence of inputs through brainstem and limbic integration. Figure 36 is the Sankey representation of the same flow architecture. Figure 37 is the network diagram emphasizing the dense interconnection of pathways.

C.4 Comparison and dose-response visualizations

The comparison and dose-response visualizations support the central convergent five-sense argument. Figure 4 shows cortisol-reduction magnitudes across interventions. Figure 7 is the additive cortisol trajectory model. Figure 19 presents the Finnish-sauna dose-response data from the Laukkanen cohort. Figure 20 shows the niacinamide and ectoin clinical dose-response curves. Figure 25 is the side-by-side single vs multisensory comparison demonstrating the roughly doubled effect sizes across outcome domains. Figure 28 is the cartogram-style sizing of sensory channels by evidence weight. Figure 39 is the gauge-meter visualization of cortisol-reduction trajectory over 12 weeks. Figure 40 is the radar chart of multimodal vs single-modal outcome dimensions. Figure 41 is the waterfall chart of cumulative cortisol reduction. Figure 42 is the adoption-and-adherence funnel. Figure 43 is the 100% stacked-bar mechanism comparison. Figure 46 is the modeled cortisol trajectory over 24 weeks comparing single-modal and multisensory protocols.

C.5 Active portfolio visualizations

The active portfolio visualizations document the topical and ingestible nutricosmetic components of the protocol. Figure 8 shows the active portfolio organized by mechanism pathway. Figure 15 is the ingredient by mechanism heat map demonstrating the convergent multi-pathway engagement. Figure 34 is the treemap of the topical portfolio. Figure 35 is the sunburst diagram of the hierarchical organization of the protocol.

C.6 Ethnographic and temporal visualizations

The ethnographic and temporal visualizations document the cross-cultural and historical context of the synthesized evidence. Figure 6 is the sense-by-sense mapping. Figure 18 is the ethnographic comparison of multisensory rituals across cultures. Figure 31 is the dot-density visualization of cited studies by publication year. Figure 44 is the timeline of key publications from 1973 to 2024. Figure 45 is the stream graph of cumulative publication output by research field.

C.7 Quality and methodology visualizations

The quality and methodology visualizations characterize the rigor of the synthesized evidence base. Figure 10 plots study quality against effect size. Figure 27 presents the U.S. regional choropleth of stress prevalence. Figure 32 is the bubble chart of cited meta-analyses by sample size and effect size. Figure 33 is the scatter plot of effect size against publication year, showing the positive trend that reflects methodological refinement and intervention-specification improvement in more recent literature.

APPENDIX D: GLOSSARY OF KEY TERMS

Allostatic load. The cumulative physiological wear-and-tear from repeated activation of stress mediators above their adaptive ranges (McEwen and Stellar 1993). **CT-afferent.** C-tactile afferent: unmyelinated low-threshold mechanoreceptor in hairy skin responding preferentially to gentle stroking at approximately 1 to 10 cm/s, projecting to the posterior insular cortex. **Cutaneous HPA axis.** The local hypothalamic-pituitary-adrenal axis homologue expressed within human skin. Slominski and colleagues (2007, 2013) demonstrated that human skin synthesizes corticotropin-releasing

hormone and possesses the enzymatic machinery for de novo cortisol biosynthesis. Cortisol. The principal glucocorticoid hormone produced by the adrenal cortex and, as detailed in Section 3, by the skin itself. Hair cortisol concentration (HCC). Cortisol measured in hair shafts, reflecting cumulative cortisol exposure over weeks to months. HCC is the preferred biomarker for chronic-stress intervention validation (Stalder 2017). Hormesis. A biphasic dose-response phenomenon in which low-dose exposure to a stressor produces adaptive benefits while higher-dose exposure produces damage. Framework articulated by Mattson and Calabrese. HPA axis. Hypothalamic-pituitary-adrenal axis. The central neuroendocrine stress-response system. Multisensory enhancement. The phenomenon in which neurons in the superior colliculus, posterior parietal cortex, and superior temporal sulcus respond to multisensory stimuli with activations exceeding the sum of unimodal responses. Neurocosmetic. Topical or ingestible products that act on the cutaneous nervous system or that modulate neuropeptide signaling in the skin. Polyvagal theory. The framework articulated by Porges (1995, 2011) distinguishing a myelinated ventral-vagal pathway from an unmyelinated dorsal-vagal pathway and the sympathetic mobilization system. Postbiotic. Bacterial metabolites and cell-wall components used topically to restore microbial-metabolite signaling at the skin surface. Pro-Hyp. Proline-hydroxyproline. The principal bioactive dipeptide produced by digestion of collagen peptides (Iwai 2005, Yazaki 2017, Ohara 2010). Stratum corneum. The outermost layer of the epidermis, composed of corneocytes embedded in a lipid matrix dominated by ceramides, cholesterol, and free fatty acids. TEWL. Transepidermal water loss. The standard biophysical measure of skin barrier integrity. Lower TEWL indicates better barrier function. Vagal tone. The activity level of the vagus nerve, measured non-invasively through high-frequency heart rate variability (HF-HRV). Affective touch. Touch that engages the C-tactile afferent pathway and produces subjective pleasantness, autonomic engagement, and oxytocin release. Distinguished from discriminative touch by its preferred velocity (1 to 10 cm/s) and its preferential representation in the posterior insular cortex. Biophilic design. The deliberate incorporation of natural elements, fractal complexity, biomorphic forms, and prospect-refuge structure into built environments, with documented effects on cortisol, autonomic function, and cognitive performance (Browning, Ryan, and Clancy 2014; Ulrich 1984). Biomarker. A measurable indicator of a biological state or process. Principal biomarkers used in this synthesis include salivary cortisol (acute), hair cortisol concentration (chronic), high-frequency heart rate variability (vagal tone), transepidermal water loss (barrier function), and Cutometer R-parameters (skin elasticity). Cohen's d. The most widely used standardized effect-size metric, defined as the difference between two means divided by the pooled standard deviation. Cohen's benchmarks: 0.2 small, 0.5 medium, 0.8 large.

Forest plot. A visualization standard in meta-analysis showing point estimates and confidence intervals for individual studies and the pooled estimate. Used throughout Section 5 to present per-channel evidence.

Hedges' g. A variant of Cohen's d that corrects for small-sample bias. Used by Packheiser et al. (2024) and other contemporary meta-analyses.

HF-HRV. High-frequency heart rate variability, measured in the 0.15 to 0.40 Hz frequency band. The principal non-invasive measurement of vagal cardiac control and ventral-vagal tone.

Hyaluronic acid synthase 2 (HAS2). The principal enzyme producing high-molecular-weight hyaluronic acid in dermal fibroblasts.

Pro-Hyp dipeptide stimulates HAS2 transcription approximately 2.3-fold (Ohara 2010), underlying the hydration benefit of marine collagen ingestion.

Lipopolysaccharide (LPS). A bacterial cell-wall component that, when it enters systemic circulation through intestinal permeability, produces systemic inflammation. The mechanism connects gut dysbiosis to skin inflammation in the gut-skin axis.

POMC. Proopiomelanocortin. The precursor protein that yields ACTH, alpha-MSH, beta-endorphin, and other peptides through tissue-specific cleavage. Expressed in the pituitary and, as documented by Slominski, in human skin.

SCORAD. Scoring Atopic Dermatitis. A composite clinical score combining extent, intensity, and subjective symptoms of atopic dermatitis. Used as the primary endpoint in ceramide and ectoin RCTs cited in Section 5.6.

Stratum corneum lipid lamellae. The multilayered lipid sheets between corneocytes that constitute the principal water-permeability barrier of the skin. Composed primarily of ceramides, cholesterol, and free fatty acids in an approximately 1:1:1 molar ratio.

Salivary cortisol. Cortisol measured in saliva, reflecting the free (non-protein-bound) bioactive fraction of circulating cortisol. The standard acute biomarker for stress-reduction intervention studies.

Substance P. A neuropeptide released by cutaneous sensory nerves under stress conditions, with documented roles in mast-cell degranulation, neurogenic inflammation, and stress-induced skin pathology.

Mast cells. Resident immune cells in the dermis that respond to CRH and substance P with degranulation, releasing histamine, tryptase, and pro-inflammatory cytokines. Central to the stress-skin connection (Theoharides and Conti 2004).

Phytoncides. Volatile organic compounds released by trees and plants with documented effects on natural killer cell function and stress markers. Underlies part of the shinrin-yoku forest-bathing effect (Li 2010).

Resonance-frequency breathing. Paced breathing at approximately 6 breaths per minute (0.1 Hz) that produces maximal amplitude of respiratory sinus arrhythmia and the largest HF-HRV elevation. The principal mechanism of HRV biofeedback (Lehrer 2013).

Stokes-Pillsbury hypothesis. The 1930 proposition that emotional states alter intestinal microbiota composition, increase intestinal permeability, and contribute to systemic inflammation that manifests dermatologically. Revived and extended by Bowe and Logan (2011) with contemporary microbiome science.

APPENDIX E: COMPLETE LIST OF FIGURES

The synthesis includes 46 figures distributed throughout the body of the paper. This appendix provides a consolidated reference list with brief descriptions and section locations for each figure.

Figure 1	The demographic case for cortisol-reduction intervention (Section 1.1)
Figure 2	Mechanistic chain linking chronic psychological stress to visible skin aging (Section 3)
Figure 3	Cortisol cascade rendered as input-mediator-outcome flow (Section 3)
Figure 4	Multisensory protocol delivered in two complementary modalities (Section 4)
Figure 5	Master forest plot of pooled effect sizes across principal modalities (Section 5)
Figure 6	Sense-by-sense mapping of in-person ritual to take-home extension (Section 4)
Figure 7	Forest plot of music interventions across outcomes (Section 5.1)
Figure 8	Forest plot of touch interventions across outcomes (Section 5.2)
Figure 9	Forest plot of olfactory and aromatherapy interventions (Section 5.4)
Figure 10	Study quality plotted against effect size (Section 5)
Figure 11	Histogram of effect-size estimates (Section 5)
Figure 12	Ingredient by mechanism heat map (active portfolio) (Section 5.6)
Figure 13	Topical portfolio organized into mechanism pathways (Section 5.6)
Figure 14	Dose-response curves for niacinamide and ectoin (Section 5.6)
Figure 15	Cross-cultural multisensory ritual comparison (Section 10)
Figure 16	Finnish sauna dose-response evidence (Section 10)
Figure 17	Five sensory channels projecting through non-overlapping afferent pathways (Section 11)
Figure 18	Modeled cortisol trajectory comparing single-modality and five-sense protocols (Section 11)
Figure 19	Cortisol-reduction magnitudes across principal interventions (Section 11)
Figure 20	Forest plot of nature exposure on cortisol (Section 5.5)
Figure 21	Forest plot of aromatherapy interventions (Section 5.4)
Figure 22	Pie chart: participants by sensory-channel meta-analysis (Section 5)
Figure 23	Donut chart: mechanism pathway distribution (Section 5)
Figure 24	Pictograph of stress prevalence in U.S. adults (Section 1)
Figure 25	Side-by-side single vs multisensory comparison (Section 11)
Figure 26	Evidence-totality dashboard (Section 1)
Figure 27	U.S. regional choropleth of stressor prevalence (Section 11)
Figure 28	Cartogram-style sensory channels resized by evidence weight (Section 11)
Figure 29	Mechanism engagement heat map (Section 6)
Figure 30	Flow map of five sensory pathways converging on integration (Section 6)
Figure 31	Dot-density visualization of cited studies by year (Section 6)
Figure 32	Bubble chart of meta-analyses by sample size and effect (Section 5)
Figure 33	Scatter plot of effect size by publication year (Section 5)
Figure 34	Treemap of topical active portfolio (Section 5.6)
Figure 35	Sunburst hierarchy of protocol mechanisms (Section 5.6)
Figure 36	Sankey diagram of channels to mechanisms to outcomes (Section 5.6)
Figure 37	Network diagram of sensory channels and sub-mechanisms (Section 7)
Figure 38	Histogram of effect sizes (extended) (Section 7)
Figure 39	Gauge meters of cortisol-reduction progress over 12 weeks (Section 7)

Figure 40	Radar chart of multisensory vs single-modality across dimensions (Section 12)
Figure 41	Waterfall chart of cumulative cortisol reduction (Section 12)
Figure 42	Funnel chart of adoption and adherence (Section 12)
Figure 43	100% stacked-bar comparison of mechanism contribution (Section 13)
Figure 44	Timeline of key publications, 1973-2024 (Section 13)
Figure 45	Stream graph of publication output by research field (Section 13)
Figure 46	Modeled cortisol trajectory over 24 weeks (Section 13)

REFERENCES

American Psychiatric Association (2024). American Adults Express Increasing Anxiousness in Annual Poll. Press release, 1 May 2024.

American Psychological Association (2023, 2024). Stress in America. Harris Poll surveys, 2023 and 2024 waves.

Antonelli M, Barbieri G, Donelli D (2019). Effects of forest bathing (shinrin-yoku) on levels of cortisol as a stress biomarker: systematic review and meta-analysis. *Int J Biometeorol*, 63(8):1117-1134.

Asserin J, Lati E, Shioya T, Prawitt J (2015). The effect of oral collagen peptide supplementation on skin moisture and the dermal collagen network. *J Cosmet Dermatol*, 14(4):291-301.

Aubert C, Chalot G (2020). Physicochemical characteristics, vitamin C, and polyphenolic composition of blood-flesh peach cultivars. *J Food Compos Anal*, 86:103337.

Belisario et al. (2023). Quantitative and qualitative determination of carotenoids and polyphenolic compounds in selected cultivars of *Prunus persica* L. PMC10039266.

Bissett DL, Miyamoto K, Sun P, Li J, Berge CA (2004). Topical niacinamide reduces yellowing, wrinkling, red blotchiness, and hyperpigmented spots in aging facial skin. *Int J Cosmet Sci*, 26(5):231-238.

Bissett DL, Oblong JE, Berge CA (2005). Niacinamide: A B Vitamin that Improves Aging Facial Skin Appearance. *Dermatol Surg*, 31(7 Pt 2):860-865.

Bolke L, Schlippe G, Gerss J, Voss W (2019). A Collagen Supplement Improves Skin Hydration, Elasticity, Roughness, and Density: Results of a Randomized, Placebo-Controlled, Blind Study. *Nutrients*, 11(10):2494.

Bowe WP, Logan AC (2011). Acne vulgaris, probiotics and the gut-brain-skin axis - back to the future? *Gut Pathogens*, 3:1.

Bowe WP, Patel NB, Logan AC (2014). Acne vulgaris, probiotics and the gut-brain-skin axis: from anecdote to translational medicine. *Benef Microbes*, 5(2):185-199.

Brooks AW et al. (2016). Don't stop believing: Rituals improve performance by decreasing anxiety. *Organ Behav Hum Decis Process*, 137:71-85.

Browning WD, Ryan CO, Clancy JO (2014). 14 Patterns of Biophilic Design. *Terrapin Bright Green*.

Byrd AL, Belkaid Y, Segre JA (2018). The human skin microbiome. *Nat Rev Microbiol*, 16(3):143-155.

Chanda ML, Levitin DJ (2013). The neurochemistry of music. *Trends Cogn Sci*, 17(4):179-193.

Choi EH, Brown BE, Crumrine D et al. (2005). Mechanisms by which psychologic stress alters cutaneous permeability barrier homeostasis and stratum corneum integrity. *J Invest Dermatol*, 124(3):587-595.

Critchley HD, Garfinkel SN (2017). Interoception and emotion. *Curr Opin Psychol*, 17:7-14.

Croy I, Bierling A, Sailer U, Ackerley R (2021). Individual Variability of Pleasantness Ratings to Stroking Touch Over Different Velocities. *Neuroscience*, 464:33-43.

Danessa G, Notario D, Regina R (2025). Effects of collagen-based supplements on skin's hydration and elasticity during ageing: A systematic review and meta-analysis. *Indian J Dermatol Venereol Leprol*, 91:730-740.

Della Loggia R, Tubaro A, Sosa S, Becker H, Saar S, Isaac O (1994). The role of triterpenoids in the topical anti-inflammatory activity of *Calendula officinalis* flowers. *Planta Med*, 60(6):516-520.

de Miranda RB, Weimer P, Rossi RC (2021). Effects of hydrolyzed collagen supplementation on skin aging: a systematic review and meta-analysis. *Int J Dermatol*, 60(12):1449-1461.

Denda M, Tsuchiya T, Elias PM, Feingold KR (2000). Stress alters cutaneous permeability barrier homeostasis. *Am J Physiol Regul Integr Comp Physiol*, 278(2):R367-R372.

Ditzen B, Neumann ID, Bodenmann G et al. (2007). Effects of different kinds of couple interaction on cortisol and heart rate responses to stress in women. *Psychoneuroendocrinology*, 32(5):565-574.

Dold M, Bartova L, Volz HP et al. (2023). Efficacy of Silexan in patients with anxiety disorders: meta-analysis of RCTs. *Eur Arch Psychiatry Clin Neurosci*, 273(7):1615-1628.

Du B, Bian Z, Xu B (2014). Skin health promotion effects of natural beta-glucan derived from cereals and microorganisms: A review. *Phytother Res*, 28(2):159-166.

Ebner F, Heller A, Rippke F, Tausch I (2002). Topical use of dexpanthenol in skin disorders. *Am J Clin Dermatol*, 3(6):427-433.

de Witte M, Spruit A, van Hooren S, Moonen X, Stams GJ (2020). Effects of music interventions on stress-related outcomes: systematic review and two meta-analyses. *Health Psychology Review*, 14(2):294-324.

de Witte M, Pinho ADS, Stams GJ, Moonen X, Bos AER, van Hooren S (2022). Music therapy for stress reduction: systematic review and meta-analysis. *Health Psychology Review*, 16(1):134-159.

Gil MI, Tomas-Barberan FA, Hess-Pierce B, Kader AA (2002). Antioxidant capacities, phenolic compounds, carotenoids, and vitamin C contents of nectarine, peach, and plum cultivars. *J Agric Food Chem*, 50(17):4976-4982.

Goel N, Kim H, Lao RP (2005). An olfactory stimulus modifies nighttime sleep in young men and women. *Chronobiology International*, 22(5):889-904.

Grice EA, Segre JA (2011). The skin microbiome. *Nat Rev Microbiol*, 9(4):244-253.

Hagerhall CM, Purcell T, Taylor R (2004). Fractal dimension of landscape silhouette outlines as a predictor of landscape preference. *J Environ Psychol*, 24(2):247-255.

Hakozaki T, Minwalla L, Zhuang J et al. (2002). The effect of niacinamide on reducing cutaneous pigmentation and suppression of melanosome transfer. *Br J Dermatol*, 147(1):20-31.

Heinrich U, Garbe B, Tronnier H (2007). In vivo assessment of ectoin: a randomized, vehicle-controlled clinical trial. *Skin Pharmacol Physiol*, 20(4):211-218.

Hobson NM, Schroeder J, Risen JL, Xygalatas D, Inzlicht M (2018). The Psychology of Rituals: An Integrative Review and Process-Based Framework. *Pers Soc Psychol Rev*, 22(3):260-284.

Holt-Lunstad J, Birmingham WA, Light KC (2008). Influence of a warm touch support enhancement intervention. *Psychosomatic Medicine*, 70(9):976-985.

Hwang E, Shin S (2015). The effects of aromatherapy on sleep improvement: systematic literature review and meta-analysis. *J Altern Complement Med*, 21(2):61-68.

Inoue N, Sugihara F, Wang X (2016). Ingestion of bioactive collagen hydrolysates enhance facial skin moisture and elasticity and reduce facial ageing signs. *J Sci Food Agric*, 96(12):4077-4081.

Iwai K, Hasegawa T, Taguchi Y et al. (2005). Identification of food-derived collagen peptides in human blood after oral ingestion of gelatin hydrolysates. *J Agric Food Chem*, 53(16):6531-6536.

Kaplan R, Kaplan S (1989). *The Experience of Nature: A Psychological Perspective*. Cambridge University Press.

Khalsa SS, Adolphs R, Cameron OG et al. (2018). Interoception and Mental Health: A Roadmap. *Biol Psychiatry Cogn Neurosci Neuroimaging*, 3(6):501-513.

Kim DU, Chung HC, Choi J, Sakai Y, Lee BY (2018). Oral Intake of Low-Molecular-Weight Collagen Peptide Improves Hydration, Elasticity, and Wrinkling in Human Skin: A Randomized, Double-Blind, Placebo-Controlled Study. *Nutrients*, 10(7):826.

Koizumi S, Inoue N, Shimizu M et al. (2018). Effects of Dietary Supplementation with Fish Scales-Derived Collagen Peptides on Skin Parameters and Condition. *Int J Pept Res Ther*, 24:397-402.

Krutmann J (2009). Pre- and probiotics for human skin. *J Dermatol Sci*, 54(1):1-5.

Lang M, Kratky J, Shaver JH, Jerotijevic D, Xygalatas D (2015). Effects of Anxiety on Spontaneous Ritualized Behavior. *Curr Biol*, 25(14):1892-1897.

Laukkanen T, Khan H, Zaccardi F, Laukkanen JA (2015). Association between sauna bathing and fatal cardiovascular and all-cause mortality events. *JAMA Intern Med*, 175(4):542-548.

Laukkanen T, Kunutsor SK, Khan H, Willeit P, Zaccardi F, Laukkanen JA (2018). Sauna bathing is associated with reduced cardiovascular mortality and improves risk prediction in men and women: a prospective cohort study. *BMC Medicine*, 16:219.

Lehrner J, Marwinski G, Lehr S et al. (2005). Ambient odors of orange and lavender reduce anxiety and improve mood in a dental office. *Physiol Behav*, 86(1-2):92-95.

Lin P, Alexander RA, Liang CH et al. (2021). Collagen formula with Djulis for improvement of skin hydration, brightness, texture, crow's feet, and collagen content. *J Cosmet Dermatol*, 20(1):188-194.

Linnemann A, Ditzen B, Strahler J, Doerr JM, Nater UM (2015). Music listening as a means of stress reduction in daily life. *Psychoneuroendocrinology*, 60:82-90.

Linnemann A, Strahler J, Nater UM (2016). The stress-reducing effect of music listening varies depending on the social context. *Psychoneuroendocrinology*, 72:97-105.

Loken LS, Wessberg J, Morrison I, McGlone F, Olausson H (2009). Coding of pleasant touch by unmyelinated afferents in humans. *Nature Neuroscience*, 12(5):547-548.

Maia Campos PMBG, Franco RSB, Kakuda L et al. (2021). Oral Supplementation with Hydrolyzed Fish Cartilage Improves the Morphological and Structural Characteristics of the Skin. *Molecules*, 26(16):4880.

Marini A, Reinelt K, Krutmann J, Bilstein A (2014). Ectoïne-Containing Cream in the Treatment of Mild to Moderate Atopic Dermatitis. *Skin Pharmacol Physiol*, 27(2):57-65.

Mattson MP (2008). Hormesis defined. *Ageing Res Rev*, 7(1):1-7.

McEwen BS (1998). Protective and damaging effects of stress mediators. *NEJM*, 338(3):171-179.

McEwen BS, Stellar E (1993). Stress and the individual: mechanisms leading to disease. *Arch Intern Med*, 153(18):2093-2101.

McGlone F, Wessberg J, Olausson H (2014). Discriminative and affective touch: sensing and feeling. *Neuron*, 82(4):737-755.

Meckfessel MH, Brandt S (2014). The structure, function, and importance of ceramides in skin. *J Am Acad Dermatol*, 71(1):177-184.

Murad S, Grove D, Lindberg KA, Reynolds G, Sivarajah A, Pinnell SR (1981). Regulation of collagen biosynthesis by ascorbic acid. *Proc Natl Acad Sci USA*, 78(5):2879-2882.

Norton MI, Gino F (2014). Rituals Alleviate Grieving for Loved Ones, Lovers, and Lotteries. *J Exp Psychol Gen*, 143(1):266-272.

Ohara H, Ichikawa S, Matsumoto H et al. (2010). Collagen-derived dipeptide, proline-hydroxyproline, stimulates cell proliferation and hyaluronic acid synthesis in cultured human dermal fibroblasts. *J Dermatol*, 37(4):330-338.

Packheiser J, Hartmann H, Fredriksen K, Gazzola V, Keyzers C, Michon F (2024). A systematic review and multivariate meta-analysis of the physical and mental health benefits of touch interventions. *Nature Human Behaviour*, 8(6):1088-1107.

Park BJ, Tsunetsugu Y, Kasetani T, Kagawa T, Miyazaki Y (2010). The physiological effects of Shinrin-yoku. *Environ Health Prev Med*, 15(1):18-26.

- Pawling R, Cannon PR, McGlone FP, Walker SC (2017).* C-tactile afferent stimulating touch carries a positive affective value. *PLoS ONE*, 12(3):e0173457.
- Perez-Sanchez A, Barrajon-Catalan E, Herranz-Lopez M, Micol V (2018).* Nutraceuticals for Skin Care: A Comprehensive Review of Human Clinical Studies. *Nutrients*, 10(4):403.
- Petrowski K, Schmalbach B, Strahler J (2021).* Acute psychobiological effects of full-spectrum light in stress recovery. *Stress*, 24(1):60-69.
- Pickart L, Margolina A (2018).* Regenerative and Protective Actions of the GHK-Cu Peptide. *Int J Mol Sci*, 19(7):1987.
- Pickart L, Vasquez-Soltero JM, Margolina A (2015).* GHK Peptide as a Natural Modulator of Multiple Cellular Pathways. *Biomed Res Int*, 2015:648108.
- Porges SW (2007).* The polyvagal perspective. *Biol Psychol*, 74(2):116-143.
- Preethi KC, Kuttan R (2009).* Wound healing activity of flower extract of *Calendula officinalis*. *J Basic Clin Physiol Pharmacol*, 20(1):73-79.
- Proksch E, Schunck M, Zague V, Segger D, Degwert J, Oesser S (2014).* Oral Intake of Specific Bioactive Collagen Peptides Reduces Skin Wrinkles and Increases Dermal Matrix Synthesis. *Skin Pharmacol Physiol*, 27(3):113-119.
- Pu SY, Huang YL, Pu CM, Kang YN, Hoang KD, Chen KH, Chen C (2023).* Effects of Oral Collagen for Skin Anti-Aging: A Systematic Review and Meta-Analysis. *Nutrients*, 15(9):2080.
- Pujos M, Chamayou-Robert C, Parat C, Bonnet M et al. (2024).* Chronic stress, cortisol and skin: a study combining in vivo and reconstructed-epidermis arms. *J Cosmet Dermatol*.
- Salem I, Ramser A, Isham N, Ghannoum MA (2018).* The Gut Microbiome as a Major Regulator of the Gut-Skin Axis. *Front Microbiol*, 9:1459.
- Salimpoor VN, Benovoy M, Larcher K, Dagher A, Zatorre RJ (2011).* Anatomically distinct dopamine release during anticipation and experience of peak emotion to music. *Nature Neuroscience*, 14(2):257-262.
- Sangsuwan W, Asawanonda P (2021).* Four-weeks daily intake of oral collagen hydrolysate results in improved skin elasticity, especially in sun-exposed areas. *J Dermatolog Treat*, 32(8):991-996.
- Schroter MA, Meyer S, Hahn MB et al. (2017).* Ectoine protects DNA from damage by ionizing radiation. *Sci Rep*, 7:15272.
- Skobowiat C, Dowdy JC, Sayre RM, Tuckey RC, Slominski A (2011).* Cutaneous HPA axis homolog: regulation by ultraviolet radiation. *Am J Physiol Endocrinol Metab*, 301(3):E484-E493.
- Slominski A, Wortsman J, Tuckey RC, Paus R (2007).* Differential expression of HPA axis homolog in the skin. *Mol Cell Endocrinol*, 265-266:143-149.
- Slominski AT, Zmijewski MA, Zbytek B, Tobin DJ, Theoharides TC, Rivier J (2013).* Key Role of CRF in the Skin Stress Response System. *Endocrine Reviews*, 34(6):827-884.
- Sowndhararajan K, Kim S (2016).* Influence of Fragrances on Human Psychophysiological Activity. *Sci Pharm*, 84(4):724-751.
- Spence C (2017).* *Gastrophysics: The New Science of Eating*. Viking.
- Stahl W, Sies H (2012).* Beta-Carotene and other carotenoids in protection from sunlight. *Am J Clin Nutr*, 96(5):1179S-1184S.
- Stalder T, Steudte-Schmiedgen S, Alexander N et al. (2017).* Stress-related and basic determinants of hair cortisol in humans: a meta-analysis. *Psychoneuroendocrinology*, 77:261-274.
- Taneja P, Olausson H, Trulsson M, Svensson P, Baad-Hansen L (2021).* Defining pleasant touch stimuli: systematic review and meta-analysis. *Psychol Res*, 85(1):20-35.
- Toda M, Morimoto K (2008).* Effect of lavender aroma on salivary endocrinological stress markers. *Arch Oral Biol*, 53(10):964-968.
- Trookman NS, Rizer RL, Ford R, Mehta R, Gotz V (2009).* Immediate and long-term clinical benefits of a topical treatment for facial lines and wrinkles. *J Clin Aesthet Dermatol*, 2(3):38-43.
- Ulrich RS (1984).* View through a window may influence recovery from surgery. *Science*, 224(4647):420-421.
- Uvnas-Moberg K, Handlin L, Petersson M (2015).* Self-soothing behaviors with particular reference to oxytocin release induced by non-noxious sensory stimulation. *Frontiers in Psychology*, 5:1529.
- Vallbo AB, Olausson H, Wessberg J (1999).* Unmyelinated afferents constitute a second system coding tactile stimuli of the human hairy skin. *J Neurophysiol*, 81(6):2753-2763.

Vaughn AR, Branum A, Sivamani RK (2016). Effects of Turmeric (*Curcuma longa*) on Skin Health: A Systematic Review. *Phytother Res*, 30(8):1243-1264.

Wager TD, Atlas LY (2015). The neuroscience of placebo effects. *Nat Rev Neurosci*, 16(7):403-418.

Watanabe E, Kuchta K, Kimura M, Rauwald HW, Kamei T, Imanishi J (2015). Effects of bergamot essential oil aromatherapy on mood states, parasympathetic nervous system activity, and salivary cortisol. *Forschende Komplementarmedizin*, 22(1):43-49.

Werkhauser N, Bilstein A, Sonnemann U (2014). Treatment of allergic rhinitis with ectoine containing nasal spray and eye drops. *J Allergy*, 2014:176597.

White DJ, de Klerk S, Woods W et al. (2016). Anti-Stress, Behavioural and Magnetoencephalography Effects of an L-Theanine-Based Nutrient Drink. *Nutrients*, 8(1):53.

Yap WS, Dolzhenko AV, Jalal Z, Hadi MA, Khan TM (2019). Efficacy and safety of lavender essential oil (Silexan) capsules among patients suffering from anxiety disorders: network meta-analysis. *Scientific Reports*, 9:18042.

Yazaki M, Ito Y, Yamada M et al. (2017). Oral Ingestion of Collagen Hydrolysate Leads to the Transportation of Highly Concentrated Gly-Pro-Hyp and Its Hydrolyzed Form of Pro-Hyp into the Bloodstream and Skin. *J Agric Food Chem*, 65(11):2315-2322.